



# **Efficacy of Certain Chemicals and Botanicals against Shoot and Fruit Borer, *Leucinodes orbonalis* (Guenee) on Brinjal (*Solanum melongena* L.)**

**Shitanshu Upadhyay <sup>a\*</sup>, Anoorag R Tayde <sup>a</sup>  
and Akhilesh Tripathi <sup>b</sup>**

<sup>a</sup> Department of Entomology, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj (U.P), India.

<sup>b</sup> Department of Entomology, KAPG college, Prayagraj, U.P, India.

## **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## **Article Information**

DOI: <https://doi.org/10.9734/ijpss/2024/v36i64696>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117317>

**Original Research Article**

**Received: 14/03/2024**  
**Accepted: 18/05/2024**  
**Published: 25/05/2024**

## **ABSTRACT**

The field trial was conducted at Central research farm, SHUATS, Prayagraj, U.P during Kharif season 2023 in randomized block design (RBD) with three replications consisting of eight treatment viz, Neem oil 2%, Chlorantraniliprole 18.5% SC, Spinosad 45%SC, Flubendiamide 480 SC, Indoxacarb 14.5% SC, Pangomia + Flubendiamide, Neem oil + Flubendiamide, and untreated Control, The result revealed that all insecticidal treatments were significantly superior to reduce the incidence of shoot and fruit damage of brinjal compared to untreated control. The plot treated with Chlorantraniliprole 18.5% SC recorded the lowest % of shoot and fruit damage 4.03% and 3.20 %

\*Corresponding author: E-mail: [shitanshu7393@gmail.com](mailto:shitanshu7393@gmail.com);

**Cite as:** Upadhyay, S., Tayde, A. R., & Tripathi, A. (2024). Efficacy of Certain Chemicals and Botanicals against Shoot and Fruit Borer, *Leucinodes orbonalis* (Guenee) on Brinjal (*Solanum melongena* L.). *International Journal of Plant & Soil Science*, 36(6), 882–886. <https://doi.org/10.9734/ijpss/2024/v36i64696>

in first and second sprays, respectively followed by spinosad 45% SC (4.43% and 4.15% respectively). The yield ranged from 85 to 225.50 q/ha in all treatments. The highest yield was recorded in Chlorantraniliprole 18.5 % SC (225q/ha) followed by spinosad 45% SC (190q/ha), Flubendiamide 480 SC (17.50q/ha), Indoxacarb 14.5 SC (160.35q/ha), Pangomia + Flubendiamide (140.33q/ha), Neem oil + Flubendiamide (132.40q/ha) and Neem oil 2% (120.50q/ha) as compared to control (85.00q/ha). When cost benefit worked out the best and most economical treatment was Chlorantraniliprole 18.5% SC (1:10.3) followed by Spinosad 45% SC (1:8.5), Flubendiamide 480 SC (1:7.5), Indoxacarb 14.5 SC (1:7.4), Pangomia + Flubendiamide (1:6.0), Neem oil + Flubendiamide (1:5.5) and Neem oil 2% (1:5.4) as compared to control (1:4.2).

**Keywords:** Cost benefit ratio; insecticides; *Leucinodes orbonalis*; *Solanum melongena*; yield.

## 1. INTRODUCTION

Brinjal (*Solanum melongena* Linnaeus) belongs to the family Solanaceae. Originated from India and now grown as a vegetable throughout the tropical, sub-tropical and warm temperate areas of the world. Brinjal is attacked by more than 70 insect pests, among the insect pest infesting brinjal, the major ones are shoot and fruit borer, *Leucinodes orbonalis* (Guen.) The larvae bore into tender shoots in the early stage resulting in drooping shoots, which are readily visible in the infested fields. At the later stage, caterpillars bore into flower buds and fruits, rendering the fruits unfit for consumption and marketing, resulting in direct yield losses. Although insecticidal control is one of the common means against the fruit borer, many of the insecticides applied are not effective in the satisfactory control of this pest.

## 2. MATERIALS AND METHODS

The trial was conducted for the management of Brinjal shoot and fruit borer using variety Pusa kranti purple Round and treatments Neem oil 2%, Chlorantraniliprole 18.5% SC, Spinosad 45% SC, Flubendiamide 480 SC, Indoxacarb 14.5% SC, Pangomia + Flubendiamide, Neem oil + Flubendiamide, were used at Central Research Farm (CRF), SHUATS, Uttar Pradesh during Kharif, 2023. The evaluation of insecticides was done in randomized block design with 3 replications. The insecticidal spray solution of desired concentration as per each treatment was freshly prepared every time at the experimental site just before the start of spraying operation. All the sprays were given during the evening hours by hand compression sprayer which was used for imposing the treatments. Two sprays were done during entire crop season when pest infestation reached economic threshold level, first spray was done on 10<sup>th</sup> September 2023

and second on 26<sup>th</sup> September 2023. The observation was recorded at 5 randomly selected plants in each treatment plot. The observations were investigated one day prior to insecticidal application and 3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> day after application for two sprays. For assessing the Brinjal fruit and shoot borer damage. The yield data in each treatment was recorded separately and subjected to statistical analysis in which cost benefit ratio in various treatments was calculated.

## 3. RESULTS OF DISCUSSION

All the insecticidal treatments were significantly superior to the untreated control in reducing the infestation of shoot damage on Brinjal after 1<sup>st</sup> insecticidal application. The % shoot damage recorded one day prior to spray ranged between 5.82 to 7.12 % (Table 1). After 3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> day of first spray lowest mean % of shoot damage was recorded in spinosad 45% SC (4.43) which was less effective than the check treatment Chlorantraniliprole 18.5% SC (4.03) followed by Flubendiamide 480 SC (4.65) Indoxacarb 14.5% SC (5.45), Pangomia + Flubendiamide (5.65), Neem oil + Flubendiamide (6.24) and the Neem oil 2% (6.40) was found to be least effective.

The data on the percent infestation of fruit borer on brinjal 3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> day after second spray revealed that all the chemical treatments were significantly superior over control. Among all the treatments lowest percent fruit infestation was recorded in spinosad 45% SC (4.15) which was less effective than the check treatment Chlorantraniliprole 18.5% SC (3.20) followed by Flubendiamide 480 SC (4.58) Indoxacarb 14.5% SC (5.33), Pangomia + Flubendiamide (5.95), Neem oil + Flubendiamide (6.46), and the Neem oil 2% (6.74) was found to be least effective. (Table 1).

**Table 1. Assessment of different insecticides and botanicals on the percent infestation, yield, CBR of shoot and fruit borer in Brinjal**

Treatment	Doses	1st Spray					2nd Spray				Yield q/ha	C:B Ratio	
		Percent shoot infestation					Percent fruit infestation						
		1DBS	3 DAS	7DAS	14 DAS	Mean	1DBS 3DAS	7DAS	14DAS	Mean			
T <sub>1</sub>	Neem oil 2%	20ml/L	6.01	6.54	6.37	6.69	6.40	6.83	6.44	6.97	6.74	120.50	1:5.4
T <sub>2</sub>	Chlorantraniliprole 18.5% SC	0.4ml/L	7.12	3.03	2.54	3.43	4.03	3.46	2.86	3.29	3.20	225.50	1:10.3
T <sub>3</sub>	Spinosad 45% SC	0.5ml/L	5.82	4.35	2.97	4.59	4.43	4.24	3.54	4.67	4.15	190.30	1:8.5
T <sub>4</sub>	Flubendiamide 480 SC	1ml/L	5.94	4.59	3.43	4.66	4.65	4.34	4.14	5.26	4.58	170.50	1:7.5
T <sub>5</sub>	Indoxacarb 14.5% SC	0.5ml/L	6.03	5.35	4.66	5.78	5.45	5.42	4.75	5.82	5.33	160.35	1:7.4
T <sub>6</sub>	Pangomia oil +Flubendiamide	40ml + 0.5ml/L	6.06	5.89	4.71	5.97	5.65	5.98	5.70	6.18	5.95	140.33	1:6.0
T <sub>7</sub>	Neem oil 2% +Flubendiamide	20ml+ 0.5ml/L	5.86	6.38	6.20	6.54	6.24	6.56	6.14	6.68	6.46	132.40	1:5.5
T <sub>8</sub>	Control		6.17	6.69	6.95	7.23	6.76	7.32	7.38	8.15	7.56	85	1:4.2
	<b>F- test</b>		NS	S	S	S	S	S	S	S	S		
	<b>S. Ed. (±)</b>		0.40	0.50	0.18	0.27	0.61	0.50	0.56	0.75	0.18		
	<b>C. D. (P = 0.05)</b>		-	1.07	0.37	0.57	1.27	1.07	1.19	1.61	1.27		

All the treatments were found to be superiorly over control on first and second spray and revealed that Chlorantraniliprole 18.5 % SC was superior in reducing the population of shoot and fruit borer the result are similar to finding made by Reddy and Kumar [1] and Saumya et al., (2023). Next most effective treatment was Spinosad 45% SC which was similar to the finding made by Mane and Kumar [2] and Chandar et al., [3]. Next effective Treatment was recorded in Flubendiamide 480 SC which was similar to the finding made by Reddy et al., [1] and Saumya and Tayde (2023) followed by Indoxacarb 14.5% SC which is similar to the finding made by Sankar and Kumar (2022) and Mane et al. [2]. Also Pangomia + Flubendiamide the result was similar to finding made by Verma et al. [4] Next effective was Neem oil + Flubendiamide which was similar to finding made by Pooja and Kumar [5]. Next least effective was T1 Neem oil 2% similar to finding made by Danish and Alexander [6].

The highest yield and cost benefit ratio was recorded in Chlorantraniliprole 18.5% SC (225.50q/ha) and (1:10.3) as respectively. The result is supported by the finding of Reddy and Kumar [1], Tripura et al., [7]. followed by Spinosad 45% SC (190.30 q/ha) and (1:8.5) which is similar to findings made by Mane and Kumar [2] and Devi et al., [8]. Flubendiamide 480 SC (170.50q/ha) and (1:7.5) in similar to findings made by Sowmya et al., [9]. Indoxacarb 14.5% SC (160.35q/ha) and (1:7.4) Sankara and kumar (2022) and Dwivedi et al. [10]. Pangomia + Flubendiamide (140 .33q/ha) and (1:6.0) are similar to findings made by Shobharani and Nandihalli [11] and Verma et al. [4]. Neem oil + Flubendiamide (132.40q/ha) and (1:5.5) similar to findings made by Patra et al. [12] Neem oil 2% (120q/ha) and (1:5.4) is similar to findings made by Sharma et al. [13], [14-17].

#### 4. CONCLUSION

The findings from the current investigation revealed that Chlorantraniliprole 18.5% SC is most effective treatment against brinjal shoot and fruit borer producing maximum yield and recorded highest cost benefit ratio compared to other treatments. While Spinosad 45% SC, Flubendiamide 480 SC, Indoxacarb 14.5% SC and Pangomia + Flubendiamide has shown average results. Neem oil + Flubendiamide and Neem oil 2% has least effectiveness in managing *Leucinodes orbonalis*. Botanicals are used in integrated pest management to avoid the

indiscriminate use of pesticides, which pollute the environment and are not very hazardous to beneficial insects.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Reddy CSTS, Kumar A. Efficacy of selected insecticides against brinjal shoot and fruit borer, *Leucinodes orbonalis* (Guenee). The Pharma Innovation Journal. 2022;11(4):1327-1330.
2. Mane PD, Kumar R. Bio-efficacy of new chemicals against shoot and fruit borer of brinjal. Journal of Entomology and Zoology Studies. 2020;8(6):1692-1693.
3. Chandar AS, Kumar A, Singh U, Kakde AA, Nawale JS, Narode MK. Efficacy of certain chemicals and biopesticides against brinjal shoot and fruit borer *Leucinodes Orbonalis* (Guenee). Journal of Entomology and Zoology Studie. 2020; 8(5):220-223.
4. Verma AP, Chandra U, Shakya A, Pratap R, Singh BS, Alam K. Efficacy of bio-insecticides and botanicals against brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee). The Pharma Innovation Journal. 2021;SP-10(12):1071-1077.
5. Pooja GS, Kumar A. Field efficacy and economics of different insecticides against brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee). The Pharma Innovation Journal. 2022;(7):4545-4548.
6. Danish M, Alexander K. Efficacy of biopesticides and chemical insecticides on brinjal shoot and fruit borer (*L.orbonalis*) during kharif season under agro climatic condition Prayagraj (UP). The Pharma Innovation Journal. 2022;Sp-11(10):710-715.
7. Tripura A, Chatterjee ML, Pande R, Patra S. Biorational management of brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee) in mid hills of Meghalaya. Journal of Entomology and Zoology Studies. 2017; 5(4):41-45.
8. Devi LL, Ghule TM, Chatterjee ML, Senapati AK. Effectiveness of biorational insecticides for the management of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee and on yield. Ecology,

- Environment and Conservation. 2015; 21(2):783-788.
9. Sowmya E, Pradeep S, Manjunatha DK, Ravulapenta S, Chethan KS. Bioefficacy of insecticides against *leucinodes orbonalis* on traditional cultivars of brinjal. Environment and Ecology. 2015;33(4): 1596-1598.
  10. Dwivedi RK, Tripathi A, Pal K, Singh DK. Effect and eco-friendly management of brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee) on brinjal; 2014.
  11. Shobharani M, Nandihalli BS. Efficacy of biorationals in the management of shoot borer, *Leucinodes orbonalis* Guenee on potato. Agricultural Science Digest. 2010; 30(4).
  12. Patra S, Thakur NS, Firake DM. Evaluation of bio-pesticides and insecticides against brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee) in Meghalaya of North-Eastern India. International Journal of Bio-resource and Stress Management. 2016; 7(5):1032-1036.
  13. Sharma JH, Tayde AR. Evaluation of bio-rational pesticides, against brinjal fruit and shoot borer, *Leucinodes orbonalis* Guen. on brinjal at Allahabad agroclimatic region. International Journal of Current Microbiology and Applied Sciences. 2017; 6(6):2049- 2054.
  14. Natekar MG, Rai S, Agnihotri NP. Bioefficacy of synthetic pyrethroids and their residues in brinjal fruit. Pestology. 1987;11(1):18-22. Pakistan Journal of Agricultural Research. 1987;23:33-39.
  15. Raja JB Rajendran, CM Pappiah. Management of brinjal shoot and fruit borer (*L. orbonalis* Guen.). Veg. Sci. 1999;26(2): 167-169.
  16. Roy G, Gazmer R, Sarkar S, Laskar N, Das G, Samanta A. Comparative bio efficacy of different insecticides against fruit and shoot borer (*Leucinodes orbonalis* Guenee) of brinjal and their effect on natural enemies. International Journal of Green Pharmacy. 2016;10(4):S257-S260.
  17. Maloth Poornachander, Anoorag R Tayde. Studies on management of fruit and shoot borer [*Leucinodes orbonalis* (Guenee)] using chemicals and neem oil in brinjal *Solanum melongena* L Poornachander, Tayde; IJPSS. 2016;34(22):1608-1613. Article no.IJPSS.91468.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/117317>