



Response of Zinc and Panchagavya on Growth and Yield of Sweet Corn (*Zea mays L. saccharata*)

Linda Golmei^{a++*} and Shikha Singh^{a#}

^a Department of Agronomy, Faculty of Agriculture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh, 211007, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i72587>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118740>

Received: 09/04/2024

Accepted: 13/06/2024

Published: 15/06/2024

Original Research Article

ABSTRACT

A field experiment was conducted during *Zaid* summer season of 2023 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj. The experiment was laid out in Randomized Block Design (RBD) with 10 treatment and replicated thrice. The results showed significantly increase in growth parameter with application of zinc (25 kg/ha) and panchagavya (3 sprays of 3%) recorded the highest plant height (166.55 cm), dry weight (69.59 g/plant) at 80 DAS. Whereas number of cobs per plant (2.00 g), number of rows per cobs (15.50 g), number of grains per cobs (28.93 g), cob yield (5.24 t/ha) were recorded at harvest. Treatment combination with zinc 25 kg/ha and panchagavya 3 sprays of 3% produced highest gross returns (Rs 2,19,560/ha), net return (Rs 1,61,952.45/ha), and benefit cost ratio (2.81) which was significantly superior to other treatments.

⁺⁺ M.Sc. Scholar;

[#] Assistant Professor;

^{*}Corresponding author: E-mail: lindagolmei700@gmail.com;

Cite as: Golmei, Linda, and Shikha Singh. 2024. "Response of Zinc and Panchagavya on Growth and Yield of Sweet Corn (*Zea Mays L. Saccharata*)". *Journal of Experimental Agriculture International* 46 (7):317-24. <https://doi.org/10.9734/jeai/2024/v46i72587>.

Keywords: Growth; panchagavya; sweet corn; yield and zinc.

1. INTRODUCTION

Maize belongs to a family poaceae is an important cereals food grain crop, also known as the “Queen of cereals” is the third most important cereals crop in India. “About two-third of the total production of maize is used for livestock feed or for commercial starch and oil production. Apart maize is an important industrial raw material and provide large opportunity for value addition” [1-4]. “Maize is cultivated in all season’s viz., Kharif, rabi and summer. In India, maize is the fifth-largest producer in the world and accounts for 3% of worldwide production, maize is grown over an area of roughly 9.18 million hectares, with a yield of 27.23 million tonnes and average productivity of 2965 kg/ha. While Uttar Pradesh gives an area of approximately 0.73 million hectares with a 7.98% to the entire country of India, which has production of approximately 1.53 million” [5].

“Sweet corn is a special type of corn, a hybridized variety of maize used as fresh or canned vegetable which produce kernels consisting mostly. The Kernel of sweet corn is much sweeter than normal. In recent years, sugar has gained more value than starch. The kernel contain a high amount of carbohydrates, protein, vitamin and minerals. Sweet corn contain 5-6% sugar, 10-11% starch, 3% water soluble polysaccharides and 70% water. The higher content of water soluble polysaccharide in the kernel adds texture and quality in addition to sweetness” [6]. “Because of its high level of glucose and low starch content sweet corn has become more common in both rural and urban areas. Being a potential crop in India sweet corn occupies important place as food, animal feed, poultry feed, industrial products mainly starch and each in brewery and seed” [7]. The crop has become increasingly important in India due to its high nutritional value and versatility in cooking.

“Among the micronutrients, zinc is one of the micronutrients recognised as an essential for the plants. Zinc plays an important role in photosynthesis, nitrogen metabolism and regulates auxin concentration in the plant. Zinc increases dry matter by increasing leaf chlorophyll content and increase in N and P efficiencies” Amanullah et al. [8]. “Zinc is crucial in taking part in plant development due to its catalytic action in metabolism for all crops especially maize. About 50% Indian soils are deficient in zinc causing low

level of zinc and yield in fodder crops and affecting the health of the livestock. Usage of the liquid manures got a lot of attention among scientists and farmer in recent years as they proved effective in increasing the absorption and translocation of nutrients by the crops. It helps to improve soil fertility and enhance crop productivity and quality of product and also working as a pest-repellent” Swaminathan et al. [9]. “Foliar application will be more efficient than soil applications at the late growth stage when there is preferential assimilate translocations into seeds (or) root activity for nutrient uptake is limited. Use of chemical fertilizers and pesticides in agriculture fields led to environmental degradation and hence as an alternative to chemicals. Panchagavya is also being sought to enhance crop establishment and health” Shakuntala et al. [10].

2. MATERIALS AND METHODS

The experiment was conducted during the *Zaid* season 2023, at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj (U.P). This experimental field is located approximately 7 kilometers from Prayagraj city, near the Yamuna River, on the left side of the Prayagraj-Rewa Road. Prayagraj is located in the subtropical zone of Uttar Pradesh, with hot summers and pleasant winters. The area’s average temperature is 22°C to 39°C. In this location, the total rainfall received during crop growing season of 2023 was 6.11 mm. The soil chemistry analysis revealed a sandy loam texture with a neutral pH of 7.4, low level of organic carbon (0.336 percent), medium level of available potassium (217.8 kg), high level of available phosphorus (28.7 kg). The soil was electrically conductive and had a conductivity of 0.618 Ds/m.

The experiment was laid out in Randomized Block Design with 10 treatments that were replicated three times viz., T₁ :120 kg N + 60 kg P + 60 kg K (control), T₂ :15 kg zn + 3 sprays of 1% Panchagavya, T₃ :15 kg zn + 3 sprays of 2% Panchagavya, T₄ :15 kg zn + 3 sprays of 3%, T₅ :20 kg zn + 3 sprays of 1% Panchagavya, T₆ :20 kg zn + 3 sprays of 2% panchagavya, T₇ :20 kg zn + 3 sprays of 3% Panchagavya, T₈ :25 kg zn + 3 sprays of 1% Panchagavya, T₉ :25 kg zn + 3 sprays of 2% Panchagavya, T₁₀ :25 kg zn + 3 sprays of 3% Panchagavya. Sweety is the variety

of sweet corn which was sown at 25 kg/ha with the spacing of 60 × 20 cm in net plot area of 3 × 3m. The field trial was ploughed, harrowed and levelled. Nitrogen was applied as per treatments and full dose of P and K. Nitrogen was applied through urea, phosphorus and applied through Di-ammonium phosphate and potassium in the form of murate of potash. Zinc was applied as basal at the time of sowing whereas the liquid foliar sprays at different time intervals as 15, 30 and 45 days after sowing (DAS). The growth parameters such as plant height (cm), dry weight per plant (g), crop growth rate and relative growth rate of data were collected from border row of plot. The number of cobs/plant, row/cob and grain/rows were collected separately from five tagged plant and green cob yield were recorded at the time of harvest from net plot area. The raw data was subjected to appropriate statistical analysis (ANOVA) as describe by Gomez and Gomez (1984). The data from the experiments were analyzed statistically, wherever treatment differences were found significant, the critical differences value were calculated at 5% level of probability (P = 0.05).

3. RESULTS AND DISCUSSION

3.1 Plant Height (cm)

Maximum plant height was observed with the application of 25 kg zinc + 3 sprays of panchagavya 3% (166.55cm) (Table 1) zinc is an activator of plant nutrients and plays an important role in growth and metabolism of microorganism Monu et al. [11]. Increase in plant height with zinc might be due cell and internodal elongation, plant metabolism, there by promoting vegetative growth which is positively correlated to the productive potentiality of plant which corroborates. Also improvement in metalloenzyme system regulatory function and growth promoting auxin production, The result was reported by Kumar and Salakinkop [12]. The application of nutrients through foliar spraying Panchagavya may have contributed to the increase in plant height because it contains a favourable combination of macro and micronutrients, growth hormones, and biofertilizers in a liquid formulation. The result was reported by Mushar et al. [13].

3.2 Dry Weight (g)

Maximum dry weight was observed with the application of 25 kg zinc + 3 sprays of panchagavya 3% (65.59 g/plant) (Table 1)

increase in dry weight which is attributed to photosynthesis being enhanced in the presence of zinc. This indicates that the micronutrients help in activating the production of tryptophan and precursor IAA, which is responsible for promoting plant growth and biomass accumulation. This result was reported by singh et al. [14]. Panchagavya make easily available of nutrients to crop without any losses (leaching, runoff), nutrient uptake will increase ultimately dry matter accumulation significantly increases was reported by Gurwinder Singh et al. [15]. The essential nutrients that organic manure provides are crucial in the plant system growth and physiological function Kumar et al. [16].

3.3 Crop Growth Rate (g/m²/day)

In the present study CGR was significantly higher for 25 kg zinc + 3 sprays of panchagavya 1% (10.19 g/g/day) (Table1) might be due to the involvement of zinc in auxin metabolism which ultimately led to hormonal activity and growth of the plant. Kumar and Bohra (2014). Application of panchagavya improved the metabolic and photosynthetic activity for enhancing biological efficiency of plant which helps the roots to spread into deeper layer of soil so that it can uptake more nutrients from soil cause there by accumulation of more carbohydrates and higher dry matter. Similar results were reported by Katyal et al [17]. The amount of all micro and macronutrients, growth enzyme present in panchagavya, ammonium, and multiplication might be the cause of the maximum dry matter [18]. Panchagavya improved synthesis of chemical substance which promote growth. Similar findings was recorded by Swaminathan et al. [9].

3.4 Yield Parameters

3.4.1 No. of cobs per plant

Maximum cobs per plant was observed with the application of 25 kg zinc + 3 sprays of panchagavya 3% (2.00) (Table 2) it is due to higher quantities of chlorophyll contents and synthesis of metabolites, oxidation and increase in activity in metabolism. Similar results were also reported by Das et al. (2020). Also it enhance the number of leaves, which increase the rate of photosynthetic and increase the quantity of assimilates enhancing the sink size. was reported by Uwah [19].

Table 1. Response of zinc and panchagavya on growth attributes of sweet corn

Treatments	Plant Height (cm) 80 DAS	Plant Dry Weight (g/plant) 80 DAS	Crop Growth Rate (g/m ² /day) 40-60 DAS
1. Control 120-60-60 kg NPK/ha	157.06	50.54	9.15
2. 15 kg Zn + 3 sprays of 1% panchagavya	157.58	53.93	9.72
3. 15 kg Zn + 3 sprays of 2% panchagavya	156.16	56.96	9.74
4. 15 kg Zn + 3 sprays of 3% panchagavya	157.49	54.74	9.28
5. 20 kg Zn + 3 sprays of 1% panchagavya	160.69	58.48	8.74
6. 20 kg Zn + 3 sprays of 2% panchagavya	160.56	57.75	9.49
7. 20 kg Zn + 3 sprays of 3% panchagavya	158.98	58.54	9.55
8. 25 kg Zn + 3 sprays of 1% panchagavya	164.72	56.48	10.19
9. 25 kg Zn + 3 sprays of 2% panchagavya	165.19	59.04	9.78
10. 25 kg Zn + 3 sprays of 3% panchagavya	166.55	65.59	9.99
SEm (+)	2.26	2.02	0.25
CD (p=0.05)	6.72	5.99	0.74

Table 2. Response of zinc and panchagavya on yield attributes of sweet corn

Treatments	No. of Cobs Per Plant	No. of Rows Per Cob	No. of Grains Per Rows	Cob Yield (t/ha)
1. Control 120-60-60 kg NPK/ha	1.80	13.26	26.60	4.44
2. 15 kg Zn + 3 sprays of 1% panchagavya	1.93	14.26	27.53	4.28
3. 15 kg Zn + 3 sprays of 2% panchagavya	1.93	14.40	26.86	4.41
4. 15 kg Zn + 3 sprays of 3% panchagavya	1.86	14.86	26.86	4.93
5. 20 kg Zn + 3 sprays of 1% panchagavya	1.86	14.00	27.26	4.73
6. 20 kg Zn + 3 sprays of 2% panchagavya	1.80	14.53	27.33	4.54
7. 20 kg Zn + 3 sprays of 3% panchagavya	1.73	14.26	26.93	4.67
8. 25 kg Zn + 3 sprays of 1% panchagavya	1.93	14.40	27.73	4.52
9. 25 kg Zn + 3 sprays of 2% panchagavya	2.00	14.73	28.00	4.92
10. 25 kg Zn + 3 sprays of 3% panchagavya	2.00	15.50	28.93	5.24
SEm (+)	0.06	0.33	0.43	0.02
CD (p=0.05)	0.16	0.98	1.28	0.06

Table 3. Response of zinc and Panchagavya on economic of sweet corn

Treatments	Cost of Cultivation (INR/ha)	Gross Return (INR/ha)	Net Return (INR/ha)	B:C Ratio
1. Control 120-60-60 kg NPK/ha	55,163.80	1,84,824	1,29,660.20	2.35
2. 15 kg Zn + 3 sprays of 1% panchagavya	56,445.05	1,82,912	1,26,235.70	2.23
3. 15 kg Zn + 3 sprays of 2% panchagavya	56,676.80	1,88,394	1,31,715.70	2.32
4. 15 kg Zn + 3 sprays of 3% panchagavya	56,907.55	2,03,314	1,46,406.45	2.57
5. 20 kg Zn + 3 sprays of 1% panchagavya	56,795.05	2,02,250	1,45,454.95	2.56
6. 20 kg Zn + 3 sprays of 2% panchagavya	57,026.30	1,90,372	1,33,345.70	2.33
7. 20 kg Zn + 3 sprays of 3% panchagavya	57,257.55	2,00,342	1,43,084.45	2.49
8. 25 kg Zn + 3 sprays of 1% panchagavya	57,145.05	1,88,744	1,31,598.95	2.30
9. 25 kg Zn + 3 sprays of 2% panchagavya	57,376.30	2,04,984	1,47,607.70	2.57
10. 25 kg Zn + 3 sprays of 3%panchagavya	57,607.55	2,19,560	1,61,952.45	2.81

3.4.2 No. of rows per cobs

Maximum rows per cobs was observed with the application of 25 kg zinc + 3 sprays of panchagavya 3% (15.5) (Table 2) the above result is due to the role of zinc photosynthesis, assimilation, and translocation of photosynthates from sources to sink in similar results was also reported by Innocent et al. [20]. Zinc has a beneficial effect on growth and physiological process resulting in higher no. of rows per cob. Soil application of zinc results in the improvement of nutrients use efficiency which leads to better cob size. Similar result were reported by Ravi et al. [21].

3.4.3 No. of grains per row

Maximum grain per rows was observed with the application of 25 kg zinc + 3 sprays of panchagavya 3% (28.93) (Table 2) zinc application give significant highest no. of grains per row because it activates several plant enzymes that are involved in carbohydrate metabolism, protein synthesis and pollen formation as reported by Ehsanullah et al. [22], Nikhil and Salakinkop (2018). The quantities of panchagavya when applied as foliar spray could have stimuli in the plant system and increased the production of growth regulator in cell system and the action of growth regulators in plant system stimulated the necessary growth and development as reported by Yadav and Loudraj (2007).

3.4.4 Cob yield (t/ha)

Maximum cob yield was observed with the application of 25 kg zinc + 3 sprays of panchagavya 3% (5.24) (Table 2) the increase in yield might be enhanced micronutrients availability. The positive effect of applied zinc on the physiological and metabolic function of the plants, might have helped to the increase in cob production process. Similar results were also reported by Ghodpage et al. [23]. The easy transfer of nutrients through foliar spray of panchagavya might be the reason for enhancement of yield attributes, and then increased yield ultimately Yadav and Loudraj [24]. Foliar application of Panchagavya at different growth stages as a growth regulator result in higher cob yield Sridhar et al. [25].

3.5 Economics

The maximum cost of cultivation (Rs 57,607.55/ha), gross return (Rs 2,19,560/ha), net

return (Rs 1,61,952.45/ha) and benefit cost ratio (B:C) (2.81) was observed with the application of 25 kg zinc + 3 sprays of panchagavya 3% (Table 3) the application of zinc nutrients result in increased growth and yield attributes of crop which ultimately increase the productivity of crop. Similar result were recorded by Swati et al. 2014 and Jagdeep et al. (2021). Foliar application of panchagavya physically supplied nutrients and growth hormone, particularly IAA and GA₃, which are found in the plant and may have stimulated the synthesis of growth regulators throughout the cell system. Similar finding was recorded by Yadav and Lourduraj [24-28].

4. CONCLUSION

The study showed that different treatment effects on almost of the growth and yield parameters. In the present investigation it was concluded that application of zinc 25 kg/ha + 3 sprays of panchagavya 3% was found to profound on growth and yield attributes in sweet corn.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Sri SRS, Debbarma V, Reddy SRK. Effect of Organic Manures and Bio-fertilizers on Growth and Yield of Baby Corn. Int. J. Plant Soil Sci. 2023;35(20):436-42. Available: <https://journalijpss.com/index.php/IJPSS/article/view/3824> [Accessed on 2024 Jun. 6].
2. Sushmitha M, Kumar BR, Jyothi PA, Gurumurthy P. Effect of Liquid Organic Manures on the Performance of Sweet Corn (*Zea mays* L. saccharate). Int. J. Environ. Clim. Change. 2023;13 (11):2628-32. Available: <https://journalijecc.com/index.php/IJECC/article/view/3431> [Accessed on 2024 Jun. 6].
3. Praburaman L, Park JH, Park YJ, He Z, Kamala-Kannan S, Oh BT. Effect of

- panchakavya (*organic formulation*) on phytoremediation of lead and zinc using *Zea mays*. Chemosphere. 2020;246:125810.
4. Aher SB, Lakaria BL, Kaleshananda S, Singh AB. Concentration and uptake of micronutrients (Fe, Zn, Cu and Mn) in soybean and wheat under organic, biodynamic and inorganic nutrient management in semi-arid tropical conditions of central India. Communications in Soil Science and Plant Analysis. 2022;53(17):2229-44.
 5. Anonymous. Agricultural Statistics at a Glance. 2019; 2020.
 6. Ventakesh S, Sanjav R, Shekhar JC. Sweet corn, speciality corn technical series I. Directorate of Maize Research, New Delhi. 2003;1-3.
 7. Das S, Jat ML, Singh KP, Rai HK. Agro – economic analysis of maize based cropping system in India. Indian Journal of Fertilizer. 2008;4(4):53-62.
 8. Amanullah S, Amir Asif Iqbal, Fahad S. Foliar phosphorus and zinc application improve growth and productivity of maize. Under moisture stress condition in semi-arid climates. Journal of Microbial & Biochemical Technology. 2016;433-439.
 9. Swaminathan C, Swaminathan V, Vijayalakshmi K. Panchagavya – Boon to organic farming, International Book, Distributing Co., Lucknow; 2007.
 10. Shakuntala NM, Vasudevan SN, Patil SB, Doddagoudar SR, Macha RCMSI, Vijaykumar AG. Organic biopriming on seed vigour inducing enzyme in paddy – An alternative to inorganics. Ecoscan. 2012;1:251-257.
 11. Monu K, Singh S, Singh V, Singh K, Richa K. Effect of zinc and boron on growth and yield of maize. Progressive research – An International Journal. 2019;14(3):215221.
 12. Kumar N, Salakinkop S. Agronomic biofortification of maize with zinc and iron micronutrients. Modern Concepts & Development in Agronomy; 2018. DOI: 10.31031/MCDA.000522.
 13. Mushar RR, Ramani BB, Lakhnotra NK. Response of summer pear millet (*Pennisetum glaucum* L.) R. Br. Emend. Stunz) to various bio-enhancers under organic farming. The Pharma Innovation Journal. 2023;12(11):426-429.
 14. Singh S, Singh V, Mishra P. Effect of NPK, boron and zinc on productivity and profitability of late sown kharif maize (*Zea mays* L.) in western Uttar Pradesh India. Ann Agric. Res. New Series. 2007;38(3): 310-313.
 15. Gurwinder S, Santosh K, Gurjagdeep SS, Ramandeep K. Effect of integrated nutrient management on yield of wheat (*Triticum aestivum* L.) under irrigated condition. International Journal of Chemical Studies. 2018;6(2):904-907.
 16. Kumar MV, Velayutham A, Kumar NS, Vasanthi D. Influence of different organic manures on the growth and yield of Baby corn. International Journal of Advances in Agricultural Science and Technology. 2018; 7(5):167-174.
 17. Katyal V, Gangwar KS, Gangwar B. Long term effect of fertilizer use on yield sustainability and soil fertility in rice-wheat system in subtropical India. Fertilizer News. 2003;48;(7):43-46.
 18. Sreethu S, Shikha S. Effect of nitrogen and Panchagavya on growth and yield of baby corn (*Zea mays* L.). The Bioscan. 2020;15 (2):243-246.
 19. Uwah DF, Ogar CC. Effect of poultry manure and plant population on soil properties and agronomic performance of sweet maize (*Zea mays* L. saccharate strut). International Journal of Applied Sciences and Technology. 2014;4(4):190-201.
 20. Innocent M, Doong Y, Wansim AN, Hai D, Etienne N, Angelique T, Hai J. Effect of combined zinc and iron application rates on summer maize yield, photosynthetic capacity and grain quality. International Journal Agronomy, Res. 2018;12(5):36-46.
 21. Ravi N, Basavarajappa R, Chandrashekar CP, Harlapur SI, Hosamani MH, Manjunatha MV. Effect of integrated nutrient management on growth and yield of quality protein maize. Journal of Agricultural Sciences. 2012;25(3):395- 395.
 22. Ehsanullah TA, Randhawa MA, Nadeem M, Naeem M. Exploring the role of Zinc in maize through soil and foliar application. Universal Journal of Agricultural Research. 2015;3(3):69-75.
 23. Ghodpage RM, Balpanda SS, Babhulkar VP, Pongade S. Effect of phosphorus and zinc fertilization on nutrient content in root, yield and nutritional quality of maize. Journal of Soil and Crops. 2008;18:458-461.
 24. Yadav BK, Lourduraj AC. Effect of organic manures and Panchagavya spray on yield

- attributes, yield and economics of rice. Crop Research. 2006;31(1):1-5.
25. Sridhar S, Arumugasamy S, Vijayalakshmi K, Balasubramanian AV. Vykshayurveda – Ayurveda for plant – A User Manual. Clarion. 2001;1-6.
26. Amutham TG, Karthikeyan R, Thavaprakash N, Barathi C. Agronomic biofortification with zinc on growth and yield of baby corn under irrigated condition. Journal of Pharmacognosy and Phytochemistry. 2019;8(3):434-437.
27. Das C, Arun K, Barik, Krishnendu Mondal. Effect of zinc application on growth and yield of baby corn (*Zea mays* L.) in lateritic soil of West Bengal. International Journal of Chemical Studies. 2020;8(2): 887-890.
28. Jagdeep S, Pratap R, Amritpal S, Naresh K, Kriti. Effect of nitrogen and zinc on growth and yield of maize. International Journal of Bio- Resource and Stress Management. 2021;12(3):179-185.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/118740>