



Effect of Light and GA₃ on Germination of Purple Passion Fruit (*Passiflora edulis* Sims) Seeds

Rivaljune Mawlieh ^{a++*}, Samir E. Topno ^{a#}
and Vijay Bahadur ^{a†}

^a Department of Horticulture, Naini Agricultural Institute, Sam Hinggibottom, University of Agriculture, Technology and Science (SHUATS), Naini, Prayagraj (Allahabad), (U.P) – 211007, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i6884>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116790>

Original Research Article

Received: 04/03/2024
Accepted: 09/05/2024
Published: 11/05/2024

ABSTRACT

The present study was carried out to investigate the effect of light and GA₃ on the germination of purple passion fruit (*Passiflora edulis* Sims) under agro climatic condition of Prayagraj. The perforated polythene bags of 4 x 5 cm size (150 gauges thickness) were used for sowing. The bags were filled with a mixture of farmyard manure (FYM), sand and soil in the ratio of 1:1:1. Pre-sowing-dip method was adopted for seed treatment. The treatments included application of [T₀: Shade net(light 20,100), T₁: GA₃ 200 ppm + light 24,500 lux, T₂: GA₃ 200 ppm + light 30,100 lux, T₃: GA₃ 250 ppm + light 24,500 lux, T₄: GA₃ 250 ppm + light 30,100 lux, T₅: GA₃ 300 ppm + light 24,500 lux, T₆: GA₃ 300 ppm + light 30,100 lux, T₇: GA₃ 350 ppm + light 24,500 lux, T₈: GA₃ 350 ppm + light 30,100 lux]. The experiment was laid in randomized block design with nine treatments and was replicated three times. Treatment T₈ performed best in term day to 1st germination(20.01), day to 50%

⁺⁺M. Sc. Scholar;

[#] Assistant Professor;

[†] Associate Professor & Head;

*Corresponding author: E-mail: mawliehreal30@gmail.com;

germination(26.68), day to complete germination(35.25),germination percentage(89.09), seed vigour index (1400.48),seedling height(10,20,30,40 days) (10.37,11.50,13.69,16.25), number of leaves (10,20,30,40 days) (3.00,4.78,6.22,9.45), shoot length (10,20,30,40 days) (4.68,4.97, 5.05,5.20), root length(10.66), shoot: root(0.62),stem girth(2.70),chlorophyll content (43.37) and survivality percentage(83.65).

Keywords: Light; GA₃; germination; seed vigour index and seedling growth; purple passion fruit.

1. INTRODUCTION

“Species within the Passifloraceae family are primarily native to regions with tropical climates. Passiflora is a large genus in the family Passifloraceae consisting of approximately 500 species, most of which are cultivated for edible fruits, pharmaceutical properties, and ornamental characteristics. Most species are herbaceous, perennial vines with a rapid growth rate. Some of them like maypop (*P. incarnata*) are considered weeds due to their rampant growth” [1]. “Passion fruit is botanically called as *Passiflora edulis* belongs to the family passifloraceae and found in tropical America (Brazil). It is known for its wonderful aroma, flavor and medicinal as well as nutritional importance. Fruits are rich in copious amount of Vitamin A, Vitamin C and mineral salts. It is commercially cultivated in countries like Kenya, Australia, New Zealand, Hawaii, South Africa and Sri Lanka” (Sema and Maiti, 2009). “The genus *Passiflora* comprising about 500 species, out of which 50 species bears edible fruits. But only two species are commercially cultivated i.e, *Passiflora edulis* var *edulis* (Purple passion fruit) and *Passiflora edulis* var *flavicarpa* (Yellow passion fruit). Some of the other important species are *P. quadrangularis*, *P. incarnata*, *P. ligularis*, *P. laurifolia* are cultivated in limited scale for local consumption” (Ramaiya et al. 2018). “The low rate of *Passiflora* seed germination may be caused by exogenous dormancy, which may be a result of both chemical and mechanical dormancy, with the latter occurring because the seeds contain inhibitors” [2]. In this regard, studies on germination of *P. ligularis* are scarce. During germination, GA₃ induces the synthesis of hydrolytic enzymes, such as amylase and protease. These enzymes degrade the stored food reserves accumulated in the endosperm or embryo as the seed matured. This degradation of carbohydrate and storage protein provides nourishment and energy to support germination and seedling growth. Gibberellins (GA₃) activate the embryonic vegetative growth, weakens the endosperm layer that involves the embryo and restricts its growth, and mobilizes the energetic

reserves from the endosperm of cereals [3,4]. Light is an extremely important factor for seed germination. However, there is disagreement in the response from different species to the absence or presence of this condition. For some species, the influence of light is favorable for germination, and for others it is negative or indifferent. This is because the light is responsible for the activation of phytochrome, a soluble chromoprotein which in the inactive form (Fv), absorbs red (R) wavelength and is transformed into an active (Fve) one [3]. Thus, this study was taken up to know the role of different concentration of GA₃ and light on seed germination of purple passion fruit.

2. MATERIALS AND METHODS

2.1 Study Location

The study was conducted at Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, located in Prayagraj, Uttar Pradesh, India. Prayagraj is located in the south-east part of Uttar Pradesh India. Prayagraj falls under agroclimatic zone IV which is named as “middle Gangetic plains” the site of experiment is located at 25.57° N latitude 81.51° E longitude and 98 meter above the sea level the temperature falls down as low as 4-5° during winter, the average rainfall in this area is around 798.900 mm annually with maximum concentration during July to September with few showers and drizzles in winter also.

2.2 Layout and Treatment Combination

The experimental was carried out to study the “Effect of light and GA₃ on seed germination of Purple Passion Fruit (*Passiflora edulis* Sims.)” in the experimental laboratory of Horticulture and field Experimentation Centre of Department of Horticulture, at the department of Horticulture, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj, (Uttar Pradesh), during 2023-24. The experimental design was RBD and there were nine treatments

which are replicated thrice. The treatment details are T₀-Shade net(light 20,100), T₁-GA₃ 200 ppm + light 24,500 lux, T₂-GA₃ 200 ppm + light 30,100 lux, T₃-GA₃ 250 ppm + light 24,500 lux, T₄-GA₃ 250 ppm + light 30,100 lux, T₅-GA₃ 300 ppm + light 24,500 lux, T₆-GA₃ 300 ppm + light 30,100 lux, T₇-GA₃ 350 ppm + light 24,500 lux and T₈-GA₃ 350 ppm + light 30,100 lux. The perforated polythene bags of 4 x 5 cm size (150 gauges thickness) were used for sowing. The bags were filled with a mixture of farmyard manure (FYM), sand and soil in the ratio of 1:1:1. Before sowing the seeds in the polythene bags, seeds were treated with GA₃. Two seeds were sown in each polythene bags. Regular watering with water-can was done gently in morning and evening. Pre-sowing dip method was adopted for seed treatment. Growth regulators of different concentrations were poured in required quantity into the labeled beakers. The observations were recorded for day to 1st germination, 50 % germination, day to complete germination, germination percentage and seed vigour index, seedling height, number of leaves, shoot length, root length, shoot: root, stem girth, chlorophyll content, survivality percentage. The experimental data collected relating to different parameters were statistically analyzed by Randomized Block Design (RBD) and results were analyzed as per the guide lines suggested by Panse and Sukhatme (1985).

3. RESULTS AND DISCUSSION

3.1 Germination Parameter

3.1.1 Day to 1st germination

The effect of light and GA₃ showed significant differences on day to 1st germination. The minimum day to 1st germination 20.01 percentage was recorded in GA₃ 350ppm + light 30,100 lux, While the maximum Day to 1st germination 38.05 percentage respectively were recorded in Shade net(light 20,100). However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux. The remarkable effect of GA₃ on minimum days required for germination might be due to it acts on the embryo and causes denova synthesis of hydrolyzing enzymes particularly amylase and protease and this hydrolyzed food is utilized for growth of embryo and thereby enhanced the germination. The effect of GA₃ on higher seed germination percentage might be due to the involvement of GA₃ in the activation of cytological enzymes along with increase in cell wall plasticity and better water absorption. These results are in agreement with the findings of

Vachhani et al. [5] in khirmi; Reshma and Simi [6] in mango.

3.1.2 Day to 50 % germination

The effect of light and GA₃ showed significant differences on day to 50 % germination. The minimum day to 50 % Germination 26.68 percentage recorded in GA₃ 350 ppm + light 30,100 lux, While the maximum 50% Germination 44.72 percentage respectively were recorded in Shade net(light 20,100). However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux.

3.1.3 Day to complete germination

The effect of light and GA₃ showed significant differences. The minimum Day to complete germination 35.25 percentage recorded in GA₃ 350 ppm + light 30,100 lux, While the maximum Day to complete germination 53.28 percentage respectively were recorded in Shade net (light 20,100). However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux.

3.1.4 Germination percentage

The effect of light and GA₃ showed significant differences on germination percentage. The maximum Germination percentage 89.69 percentage recorded in GA₃ 350 ppm + light 30,100 lux, While the minimum Germination percentage 68.66 percentage respectively were recorded in Shade net(light 20,100). However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux. GA₃ induces the denovo synthesis of proteolytic enzymes like α -amylase and ribonuclease. Amylases in turn hydrolyse starch in the endosperm, providing the essential sugars for the initiation of growth processes. GA₃ treatment is also known to overrule the photo dormancy, thermo-dormancy, dormancy imposed by incomplete embryo development, mechanical barriers, and presence of germination inhibitors. The result was in agreement with findings of Praveen et al. [7] in custard apple and Muralidhara et al. [8] in mango Lalitha et al. [9].

3.2 Growth Parameters

3.2.1 Seedling Height (cm)

The effect of light and GA₃ showed significant differences on seedling height (cm) at 10, 20, 30 and 40 days after germination of Purple Passion Fruit (*Passiflora edulis* Sims). The maximum seedling height was observed in GA₃ 350 ppm +

light 30,100 lux with (10.37, 11.50, 13.69 and 16.25). While the minimum seedling height (5.60, 6.34, 8.02 and 10.22) respectively were observed in Shade net(light 20,100) at 10, 20, 30 and 40 days after germination. However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux. Aatla and Srihari, [10] reported that treatment of mango cv. Alphonso kernel with GA₃ 500 ppm resulted in greater seedling height and internodal length. Increased shoots length with GA₃ treatment might be because it activated amylase, which digested the available carbohydrate into simple sugars so that energy and nutrition were easily available to faster growth of seedling Aatla and Srihari, [10]. GA₃ increased cell division and cell growth apparently leading to increased development of young leaves. Therefore, from the present study it is evident that the maximum number of leaves per seedling with GA₃ may be attributed to increase the physiological process and stimulatory action of GA₃ to form new leaves at faster rate. Archana et al. [11] reported similar results in *Annona squamosa*, Ramchandra and Govind, in guava and Dinesh and Padmapriya [12].

3.2.2 Number of leaves

The effect of light and GA₃ showed significant differences on number of leaves per seedling at 10, 20, 30 and 40 days after germination of Purple Passion Fruit (*Passiflora edulis* Sims). The maximum number of leaves per seedling was observed in GA₃ 350 ppm + light 30,100 lux with (3.00, 4.78, 6.22 and 9.45). While the minimum number of leaves per seedling (2.00, 2.56, 3.56 and 5.78) respectively were observed in Shade net(light 20,100) at 10, 20, 30 and 40 days after germination. However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux. GA₃ pre-soaking can be correlated with higher seed germination, higher shoot length and root length and number of leaves has lead to over all assimilation and distribution of food material with the plant [13] and hence resulted in higher seedling vigour [14]. The results were in close agreement with the Pampanna and Sulikeri (2001) in sapota; Kumar et al. [15], Padma Lay et al. [16] and Amit Desai et al. [17] in papaya.

3.2.3 Shoot length (cm)

The effect of light and GA₃ showed significant differences on shoot length (cm) at 10, 20, 30 and 40 days after germination of Purple Passion Fruit (*Passiflora edulis* Sims). The maximum

shoot length (cm) was observed in GA₃ 350 ppm + light 30,100 lux with (4.68, 4.97, 5.05 and 5.21). While the minimum shoot length (cm) (3.56, 4.01, 4.23 and 4.30) respectively were observed in Shade net(light 20,100) at 10, 20, 30 and 40 days after germination. However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux.

3.2.4 Root length (cm)

The effect of light and GA₃ showed significant differences on root length (cm) of Purple Passion Fruit (*Passiflora edulis* Sims) seeds. The maximum root length (10.66) was recorded in T₈ GA₃ 350 ppm + light 30,100 lux. whereas the minimum Root length (6.97) was found recorded in Shade net(light 20,100) respectively. However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux.

3.2.5 Seed vigour index

The effect of light and GA₃ showed significant differences on seed vigour index. The maximum seed vigour index (1400.48) was recorded in T₈ GA₃ 350 ppm + light 30,100 lux. Where as the minimum seed vigour index (701.93) was found recorded in Shade net(light 20,100) . However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux.

3.2.6 Shoot: root

The effect of light and GA₃ showed significant differences on shoot : root of Purple Passion Fruit (*Passiflora edulis* Sims) seeds. The maximum shoot : root (0.62) was recorded in T₈ GA₃ 350 ppm + light 30,100 lux. whereas the minimum shoot : root (0.49) was found recorded in Shade net(light 20,100) respectively. However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux. Ravishankaret al. [18] also reported the similar results.

3.2.7 Stem girth (mm)

The effect of light and GA₃ showed significant differences on stem girth (mm) of Purple Passion Fruit (*Passiflora edulis* Sims) seeds. The maximum stem girth (mm) (2.70) was recorded in T₈ GA₃ 350 ppm + light 30,100 lux. Where as the minimum stem girth (mm) (1.86) was found recorded in Shade net(light 20,100) respectively. However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light

30,100 lux. Choudhary et al. [19] and Kumawat et al. [20] also reported the similar results.

3.2.8 Chlorophyll content (spad)

The effect of light and GA₃ showed significant differences on chlorophyll content of Purple Passion Fruit (*Passiflora edulis Sims*) seeds. The maximum Chlorophyll content (43.37) was recorded in T₈ GA₃ 350 ppm + light 30,100 lux. Where as the minimum Chlorophyll content (25.95) was found recorded in Shade net(light 20,100) respectively. However, GA₃ 350 ppm + light 24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux. These findings are in agreement with the results obtained by Lay et al. [21] in papaya. The different levels of GA₃ did not affect significantly to increase chlorophyll content.

3.3 Survival and Establishment

3.3.1 Survivability percentage

The effect of light and GA₃ showed significant differences on survivability percentage of Purple Passion Fruit (*Passiflora edulis Sims*) seeds. The maximum Survivability percentage (83.65) was recorded in T₈ GA₃350 ppm + light 30,100 lux. Where as the minimum Survivability % (32.87) was found recorded in Shade net(light 20,100) respectively. However, GA₃ 350 ppm + light

24,500 lux are found statistically at par to GA₃ 350 ppm + light 30,100 lux. The application of GA₃ might have boosted the early germination and growth of seedling by cell multiplication and cell elongation subjected to the better plant growth resulting higher rate of seedling survival. This might be due to rapid and early germination which resulted in giving more periods for vegetative growth for better establishment of plants [22]. The combination of growing media and GA₃ significantly promoted the germination and growth by multiplication and elongation of plant cell. On the other hand, the improved soil condition, soil health, availability of the nutrients, might be supported to appropriate cation exchange capacity and water holding capacity. Thus permitting adequate moisture exchange of gases uptake of nutrients facilitate better and healthy growth resulting maximum survival of seedling under the treatments [23].

3.3.2 Economics

Due to the increase in germination percentage and survivability percentage as showed in (Table 1 and 3), the used of GA₃350 ppm + light 30,100 lux (T₈) for the preparation of passion fruit seedling have proved to be profitable and showed maximum net return (Rs. 13337.50/1000 seedlings) and B:C (2.76) for the experiment [24,25].

Table 1. Effect of light and GA₃ on Germination Parameters and seed vigour index of purple Passion fruit (*Passiflora edulis sims*)

Notion	Treatment details	Germination Parameters				Seed vigour index
		Day to 1 st germination	50 % Germination	Day to complete germination	Germination percentage	
T ₀	Shade net(light 20,100)	38.05	44.72	53.28	68.66	701.93
T ₁	GA ₃ 200 ppm + light 24,500 lux	31.32	41.84	50.40	71.76	824.51
T ₂	GA ₃ 200 ppm + light 30,100 lux	26.09	32.76	41.33	74.56	797.49
T ₃	GA ₃ 250 ppm + light 24,500 lux	31.25	37.92	46.48	74.25	820.22
T ₄	GA ₃ 250 ppm + light 30,100 lux	28.95	35.61	44.18	78.64	1079.78
T ₅	GA ₃ 300 ppm + light 24,500 lux	26.81	33.48	42.04	78.64	1198.02
T ₆	GA ₃ 300 ppm + light 30,100 lux	24.64	37.99	46.56	80.59	1069.10
T ₇	GA ₃ 350 ppm + light 24,500 lux	31.30	35.17	39.87	84.08	1366.08
T ₈	GA ₃ 350 ppm + light 30,100 lux	20.01	26.68	35.25	89.09	1400.48

Notion	Treatment details	Germination Parameters				Seed vigour index
		Day to 1 st germination	50 % Germination	Day to complete germination	Germination percentage	
F Test		S	S	S	S	S
S.Ed(±)		1.48	1.48	2.45	3.46	68.68
CV		5.98	5.28	6.76	5.45	246.98
CD at 0.5%		3.14	3.14	5.19	7.34	136.98

Table 2. Effect of light and GA₃ on seedling growth parameters of purple Passion fruit (*Passiflora edulis sims*)

Notion	Treatment	Seedling height (cm) (Day after germination)				Number of leaves per seedling (Day after germination)				Shoot length (cm) (Day after germination)			
		10	20	30	40	10	20	30	40	10	20	30	40
		Days	Days	Days	Days	Days	Days	Days	Days	Days	Days	Days	Days
T ₀	Shade net(light 20,100)	5.60	6.34	6.83	10.22	2.00	2.56	3.56	5.78	3.56	4.01	4.23	4.30
T ₁	GA ₃ 200 ppm + light 24,500 lux	5.72	6.70	8.02	11.49	2.11	3.11	4.00	6.44	3.76	4.11	4.59	4.63
T ₂	GA ₃ 200 ppm + light 30,100 lux	5.83	7.14	7.58	10.70	2.44	3.11	4.00	6.22	3.68	4.53	4.58	4.84
T ₃	GA ₃ 250 ppm + light 24,500 lux	5.95	7.19	8.08	11.05	2.22	3.22	4.00	6.78	3.82	4.71	4.78	5.05
T ₄	GA ₃ 250 ppm + light 30,100 lux	7.24	9.19	9.70	13.73	2.11	4.00	5.11	6.11	3.67	4.60	4.64	5.07
T ₅	GA ₃ 300 ppm + light 24,500 lux	7.28	9.46	9.50	15.23	2.11	3.11	5.11	8.78	3.88	4.49	4.97	5.08
T ₆	GA ₃ 300 ppm + light 30,100 lux	8.14	8.49	11.29	13.27	2.45	4.11	5.89	8.33	3.97	4.59	4.98	5.19
T ₇	GA ₃ 350 ppm + light 24,500 lux	9.10	10.63	11.39	15.72	2.67	4.45	6.00	8.78	4.26	4.95	5.04	5.20
T ₈	GA ₃ 350 ppm + light 30,100 lux	10.37	11.50	13.69	16.25	3.00	4.78	6.22	9.45	4.68	4.97	5.05	5.21
F Test		S	S	S	S	NS	S	S	S	S	S	S	S
S.Ed(±)		0.72	1.90	1.06	1.17	0.22	0.24	0.32	0.55	0.37	0.39	0.53	0.31
CV		12.11	25.05	14.93	10.94	11.86	8.24	8.03	9.11	11.71	10.41	13.71	7.83
CD at 5%		1.53	4.03	2.26	2.48	0.47	0.51	0.68	1.17	0.79	0.82	1.13	0.66

Table 3. Effect of light and GA₃ on root length, shoot:root, stem girth(mm), chlorophyll content and Survivality % of purple Passion fruit (*Passiflora edulis sims*)

Notion	Treatment details	Root length (cm)	Shoot: root	Stem girth (mm)	Chlorophyll content	Survivality %
T ₀	Shade net(light 20,100)	6.97	0.49	1.86	25.95	32.87
T ₁	GA ₃ 200 ppm + light 24,500 lux	7.60	0.54	2.00	28.4	45.98
T ₂	GA ₃ 200 ppm + light 30,100 lux	7.90	0.61	2.16	32.58	53.59
T ₃	GA ₃ 250 ppm + light 24,500 lux	8.20	0.58	2.19	32.44	61.84
T ₄	GA ₃ 250 ppm + light 30,100 lux	8.50	0.60	2.26	31.63	63.19
T ₅	GA ₃ 300 ppm + light 24,500 lux	8.90	0.57	2.43	35.16	71.49
T ₆	GA ₃ 300 ppm + light 30,100 lux	9.50	0.55	2.54	35.29	81.59
T ₇	GA ₃ 350 ppm + light 24,500 lux	9.60	0.54	2.66	42.3	82.84
T ₈	GA ₃ 350 ppm + light 30,100 lux	10.66	0.62	2.70	43.37	83.65
F Test		S	S	S	S	S
S.Ed(±)		0.548	0.012	0.032	4.73	2.345
CV		7.756	0.48	0.421	16.96	9.858
CD at 0.5%		1.161	0.023	0.065	10.02	4.681

4. CONCLUSION

On the basis of our experimental finding, it can be concluded that the best result was found in treatment T₈ (GA₃ 350 ppm + light 30,100 lux) in term day to 1st germination(20.01), day to 50% germination (26.68),day to complete germination (35.25), germination percentage (89.09), seed vigour index(1400.48),seedling height (10,20,30,40 days) (10.37,11.50,13.69,16.25), number of leaves (10,20,30,40 days) (3.00, 4.78,6.22,9.45), shootlength (10,20,30,40 days) (4.68,4.97,5.05,5.20), rootlength (10.66), shoot:root (0.62), stemgirth (2.70), chlorophyll content (43.37) and survivality percentage (83.65).

The highest B:C ratio was found in T₈ (GA₃ 350 ppm + light 30100 lux) with 2.76.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Amir Rezazadeh and Eric T. Stafne. Comparison of Seed Treatments on the Germination of Seven Passion Fruit Species. Int. J. Curr. Microbiol. App. Sci. 2018;7(11):3074-3083.
2. Delanoy M, Van Damme P,
3. Scheldeman X, Beltran J. Germination of *Passiflora mollissima* (Kunth) L. H. Bailey, *Passiflora tricuspidata* Mast. and *Passiflora nov sp.* Seeds. *Scientia Horticulturae*, Amsterdam. 2006;110:198-203, 2006.
4. Taiz L, Zeiger E. *Fisiologia vegetal*. 5 ed. Porto Alegre: Artmed. 2013:918.
5. Hota Satya Narayan, Kumar Ajay Karna, PK Jain, Bharat Dakhad. Effect of gibberellic acid on germination, growth and survival of jamun (*Syzygium cumini* L. Skeels). The Pharma Innovation Journal. 2018;7(8):323-326
6. Vachhani KB, Gohil JH, Pandey R, Ray NR. Influence of chemicals, PGRs and cow dung slurry as seed treatment on germinability, growth and development of khirni (*Manilkara hexandra* L) under net house condition. Trends Biosci. 2014;7(14):1641-164.
7. Reshma UR, Simi S. Effect of pre-sowing treatments, sowing positions and age of stones after extraction on germination of mango. Int. J Curr. Microbiol. App. Sci. 2019; 8(4):2565-2573.
8. Praveen J, Reddy BS, Patil RT. Effect of different seed treatments on germination of *Annona squamosa* L. under misthouse ecosystem, J. Asian. Hort. 2006;2(1/2): 1923.
9. Muralidhara BM, Reddy YTN, Akshitha HJ, Srilatha V. Effect of presowing treatments on germination, growth and vigour of polyembryonic mango seedlings. Environ. Ecol. 2015;33(3):1014-1018.
10. Lalitha KR, Tank RV, Chawla SL, Jena Suchismita. Effect of chemicals on seed germination and seedling growth of Aonla (*Embllica officinalis* Gaertn.). The Pharma Innovation Journal. 2020;9(12):239-243.

11. Aaltla HB, Srihari D. Influence of pre-sowing treatments on germination, growth and vigour of mango cv. Alphonso. Asian J Hort. 2013;8(1):122-125.
12. Archana C, Jadhav YL, Bhagure, Raundal RM. Effect of PGR, chemicals and plant extract on seed germination and seedling growth of custard apple (*Annona squamosa*). Asian J Hort. 2015;10(1):184-186
13. Dinesh A, Padmapriya S. Effect of different pre-sowing treatments on seed germination, seed coat morphology and survival of guava (*Psidium guajava* L.). Journal of Pharmacognosy and Phytochemistry. 2022;11(3):265-269
14. Brain PW, Hemming HG. The effects of GA on shoot growth of pea seedling. Physiology of plant, 1955;8:669681.
15. Pampanna Y, Sulikeri GS. Effect of growth regulators on seed germination and seedling growth of Sapota. Karnataka J Agric. Sci 2001;14:1030-1036.
16. Kumar P, Sehrawat SK, Dahiya OS, Dahiya DS. Papaya seed priming in relation to seed vigour. Haryana J. Hort. Sci 2011;40(1, 2):4-9.
17. Padma Lay, Basvaraju GV, Sarika G, Amrutha N. (2013). Effect of seed treatment to enhance seed quality of papaya (*Carica papaya* L.) cv. Surya. G.J.B.A.H.S 2013;2(3):221-225.
18. Desai Amit, Ashwin Trivedi, Bharat Panchal, Velji Desai. Improvement of papaya seed germination by different growth regulator and growing media under net house condition. International Journal of Current Microbiology and Applied Sciences. 2017;6(9):828-834.
19. Ravishankar Lanjhiyana, GD Sahu, HK Panigrahi and Pratibha Katiyar. Role of pre-sowing seed treatment on germination behavior and seedling vigour of papaya (*Carica papaya* L.). Journal of Pharmacognosy and Phytochemistry. 2020;9(5): 3039-3042.
20. Choudhary RC, Kanwar J, Chouhan GS, Sing P, Tanwar DR. Effect of GA3 and growing media on seedling growth of papaya (*Carica papaya* L.) cv. Pusa Nanha. Inter. J. of Che. Stud. 2018; 6(6):1008-1012.
21. Kumawat R, Maji S, Govind, Meena DC. Studies on seed germination and seedling growth of papaya (*Carica papaya* L.) cv. Coorg Honey Dew as influenced by media and chemicals. J. Crop and Weed. 2014;10(2): 281-286.
22. Lay P, Basvaraju GV, Pashte VV, Gowri M. Studies on effect of giberellic acid (GA3) and potassium nitrate (KNO3) on breaking of seed dormancy of papaya (*Carica papaya* L.) cv. Surya. The Ecoscan. 2015;9(1&2):111115.
23. Sharma RK. Effect of salicylic acid and gibberellic acid on seed germination and growth of pea. Int. J. of Pl. Sci., 2012;7(2):322-3.
24. Parmar Mohni, Pandey SK, Pandey CS, Singh Pawandeep, Ravi Aiswarya. Effect of Growing Media and GA3 on Seed Germination and Seedling Growth of Acid Lime (*Citrus aurantifolia* Swingle) Cv. Vikram. Int. J. Curr. Microbiol. App. Sci. 2019; 8(08):1260-1271.
25. Akhil Komaravalli, Bahadur Vijay, Annjoe V Joseph and Reena Shirle Lawrence. Effect of gibberellic acid and naphthalene acetic acid on germination and seedling growth on papaya (*Carica papaya* L.). The Pharma Innovation Journal. 2023;12(12): 2510-2514.
26. Anburani A, Shakila A. Influence of seed treatment on the enhancement of germination and seedling vigour of papaya. Acta Horticulturae. 2010;851:295-298.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/116790>