



Assessing the Impact of Casing Material Thickness on Yield and Biological Efficiency of Milky Mushroom (*Calocybe indica* P&C)

Marimuthu Elangovan ^{a,b}, Vaithiyalingam Gopinath ^c,
Ganesan Rathinakumar ^d, Madhaiyan Sakthivel ^e
and Mohan Kumar Biswas ^{b*}

^a ICAR-Indian Agricultural Research Institute, New Delhi- 110 012, India.

^b Palli Siksha Bhavana, Visva-Bharati, West Bengal- 721 236, India.

^c Dr.Rajendra Prasad Central Agricultural University, Bihar- 848 125, India.

^d Sethubaskara Agriculture College and Research Foundation, Tamil Nadu- 630 311, India.

^e G.B. Pant University of Agriculture and Technology, Uttarakhand- 263 145, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i101478>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124292>

Original Research Article

Received: 25/07/2024

Accepted: 27/09/2024

Published: 30/09/2024

ABSTRACT

Calocybe indica, known as the milky mushroom, has significant potential for agricultural expansion and economic development, particularly in West Bengal, India. This study investigates the impact of different casing material thicknesses (0.5", 0.75", 1.0", and 1.5") on the yield and biological

*Corresponding author: E-mail: elangomarimuthu96@gmail.com;

Cite as: Elangovan, Marimuthu, Vaithiyalingam Gopinath, Ganesan Rathinakumar, Madhaiyan Sakthivel, and Mohan Kumar Biswas. 2024. "Assessing the Impact of Casing Material Thickness on Yield and Biological Efficiency of Milky Mushroom (*Calocybe Indica* P&C)". *Journal of Advances in Biology & Biotechnology* 27 (10):556-63. <https://doi.org/10.9734/jabb/2024/v27i101478>.

efficiency of *C. indica*. Using a Completely Randomized Design (CRD), the effects on primordia initiation, sporophore maturation, yield, and biological efficiency were assessed. Results indicated that a casing thickness of 1.0" was optimal, leading to the quickest primordia initiation (10.4 days) and sporophore maturation (9 days), and yielding the highest total yield (717.5 g) and biological efficiency (71.75%). In contrast, thinner and thicker casings resulted in lower yields and efficiencies. The findings suggest that a 1.0" casing thickness significantly enhances *C. indica* cultivation, providing practical recommendations for commercial mushroom farming. This optimization can support the development of milky mushroom farming in West Bengal, offering new opportunities for local farmers and contributing to regional agricultural diversification and economic growth.

Keywords: *Calocybe indica*; casing thickness; yield; biological efficiency; commercial cultivation.

1. INTRODUCTION

The milky mushroom, scientifically known as *Calocybe indica*, is characterized by its robust and fleshy nature, with a distinctive milky white colour. It is notable for its long shelf life, which extends from 4 to 5 days, making it a viable option for both fresh consumption and extended use in culinary purposes. The discovery of this mushroom dates back to 1974, when Purkayastha and Chandra identified it in the northeastern region of India, specifically in West Bengal [1]. This initial discovery was significant but did not immediately lead to widespread cultivation or commercialization.

The importance of *Calocybe indica* gained renewed attention in 1997 when Krishnamoorthy rediscovered the mushroom in Tamil Nadu, southern India. Despite its initial discovery and subsequent rediscovery, *Calocybe indica* did not enter commercial cultivation until 1998. This year marked a pivotal moment when Krishnamoorthy and colleagues successfully cultivated the mushroom at the Regional Research Station in Aruppukkottai. This effort resulted in the introduction of a new variety, *Calocybe indica* var. APK2, which represented the first commercial cultivation of this species globally [2].

The genus *Calocybe*, which originated in India, currently encompasses nearly 40 species. These species are predominantly cultivated in tropical and subtropical regions that offer the high temperatures and relative humidity essential for their growth and development [3]. The cultivation of *Calocybe indica* has demonstrated significant potential due to its favourable nutritional profile, including high levels of protein, lipids, fiber, polysaccharides, and essential amino acids, while being low in fatty acids [4,5].

Calocybe indica primarily thrives in the southern part of the Indian subcontinent. Regions such as

Tamil Nadu, Andhra Pradesh, Karnataka, and Kerala provide the optimal climatic conditions characterized by high and moderate temperatures throughout the year that are necessary for the successful cultivation of this mushroom [3]. The climatic conditions in Orissa are also suitable for the growth of *Calocybe indica* except during the cooler months from November to February, which may impede its cultivation [6].

However, despite the favourable conditions in the northeastern part of India, including West Bengal, *Calocybe indica* has not been commercially cultivated in this region. West Bengal is predominantly agricultural, with over 90% of its cultivators being small and marginal farmers [7]. This contrasts sharply with the successful commercialization of other mushroom varieties, such as button and oyster mushrooms, which are cultivated throughout the year in West Bengal. The absence of standardized cultivation practices for milky mushrooms in this region represents a significant gap in agricultural development.

The potential for *Calocybe indica* to contribute to local agriculture and improve the livelihoods of small and marginal farmers in West Bengal is substantial. The application of a soil-sand-farmyard manure mixture on the surface of a fully colonized mushroom bag is known as casing. Casing materials are crucial in mushroom cultivation as they influence the moisture retention, temperature regulation, and overall growth conditions for the fungi. This casing layer not only provides moisture for growth and development but also offers physical support for the formation of fruiting bodies. The makeup of the casing mixture and thickness influences its quality, including factors like compactness, texture, structure, pH, water retention, and carbon-to-nitrogen ratio. These elements directly

impact mycelial growth in the casing layer and the initiation of fruiting bodies. The current study aims to exploring the effects of different casing material thicknesses on the yield and biological efficiency of *Calocybe indica* in the lateritic belt of West Bengal. By establishing optimal casing thicknesses, the study aims to provide practical recommendations for commercial cultivation, thereby enhancing the viability of milky mushroom farming in West Bengal [8].

2. MATERIALS AND METHODS

2.1 Source of Culture

The pure culture of *Calocybe indica* used in this study was obtained from the Mushroom Research Laboratory, Department of Plant Pathology, Orissa University of Agriculture and Technology, Bhubaneswar, Orissa. This culture was transferred to sterilized Potato Dextrose Agar (PDA) slants and maintained at room temperature throughout the research period.

2.2 Preparation of Mother Spawn

Healthy wheat grains were used as the substrate for the preparation of mother spawn. The grains were boiled for 30 minutes and then shade-dried for 2-3 hours. Subsequently, the grains were mixed with 2% calcium carbonate (CaCO₃) and 2% gypsum (CaSO₄) on a dry weight basis to achieve a moisture content of 50-55%. The prepared grains were placed in 250 ml conical flasks, which were then plugged with non-absorbent cotton and autoclaved at 121°C and 15 psi pressure for 2 hours. After autoclaving, the flasks were allowed to cool overnight before being inoculated aseptically with the pure culture. The inoculated flasks were then incubated at 27°C ± 2°C for 10-15 days.

2.3 Preparation of Bed Spawn

For the preparation of bed spawn, the procedure was similar to that of the mother spawn. Healthy wheat grains were prepared and mixed with calcium carbonate and gypsum, then filled into 12×18 cm clear transparent polythene bags at a rate of 150 g per bag. The bags were plugged with non-absorbent cotton and secured with rubber bands. The filled bags were autoclaved for 2 hours at 15 psi and 121°C. After autoclaving, the bags were inoculated aseptically with mother spawn and incubated in a dark place at 28-30°C for 15-20 days.

2.4 Preparation of Mushroom Beds

Substrate with 60% moisture content was filled into clear transparent polypropylene bags of size 16"×18" and 150-gauge thickness. The substrate was layered with the bottom and top layers being thinner compared to the intermediate layers. The beds were prepared with 4% spawning, employing a layering method with 4-5 layers of spawn. The mouths of the spawned beds were closed with rubber bands, and 15-20 holes were made on the surface of each bed. The beds were then suspended on nylon ropes in a dark room with an appropriate temperature range of 25-35°C and relative humidity of 80-90% to facilitate spawn run. After complete colonization of the substrate by the mushroom mycelium, casing was applied.

2.5 Preparation of Casing Material and Casing of Beds

Casing material was prepared by mixing garden soil, farmyard manure (FYM), and sand in a 1:1:1 ratio. The mixture was sterilized in an autoclave at 121°C and 15 psi for 2 hours and allowed to cool to room temperature overnight. Once the milky white mycelium of *C. indica* had fully colonized the substrate, indicating the completion of spawn run, the mouths of the mushroom beds were opened. A casing mixture with varying thicknesses of 0.5", 0.75", 1.0", and 1.5" was spread over the surface of the beds. After casing, the beds were transferred to the cropping room for primordial initiation and sporophore production. To maintain surface wetness, the beds were watered twice or thrice daily. Additionally, the cropping room's relative humidity was regulated by covering the ventilated cemented floor with sand and watering it as needed.

2.6 Harvesting of Mushrooms

Mushrooms were harvested by hand-picking, involving slight twisting and pulling of the matured fruiting bodies. During the cropping period of 50-60 days, three harvests were conducted at 10-15 days intervals. The weight of each individual fruiting body was recorded immediately using a single-pan weighing balance with 1 g sensitivity. The total yield of mushrooms from each replication was calculated by summing the fresh weights from all harvests, expressed as weight (g) per unit of dry weight substrate. Yield and biological efficiency were calculated using the following formulas:

$$\text{Yield (g)} = \frac{\text{Sum of weight of individual sporophores (g)}}{\text{Number of sporophore}}$$

$$\text{Biological efficiency (\%)} = \frac{\text{Weight of fresh mushroom (g)}}{\text{Weight of dry substrate (g)}} \times 100$$

2.7 Statistical Analysis

Data collected from the experiments were analysed using a Completely Randomized Design (CRD) methodology. The critical difference (CD) was computed at a 1% significance level to determine the significance of the results.

3. RESULTS AND DISCUSSION

The study assessed the impact of different casing layer thicknesses (0.5", 0.75", 1.0", and 1.5") on several factors, including the initiation of primordia, the maturation period of sporophores, yield, and biological efficiency of *Calocybe indica*. The results of these evaluations are detailed in Tables 1 and 2.

3.1 Time for Primordia Initiation and Sporophore Maturation

The results revealed that a casing thickness of 1.0" was optimal for both early primordia initiation (10.4 days) and sporophore maturation (9 days). This was followed by the 0.75" casing thickness, which required 14.20 days for primordia initiation and 12.20 days for sporophore maturation. The

1.5" casing thickness took 16.40 days for primordia initiation and 12.40 days for maturation. The casing thickness of 0.5" resulted in the longest periods, with 17.4 days for primordia initiation and 14.40 days for fruiting body maturation (Fig. 1). The poor mycelial growth and colonization in lower thickness of casing material may be the reason for the prolonged duration before primordia initiation occurs. The prolonged maturation time for the sporophores may be attributed to insufficient casing thickness, which fails to retain moisture for an extended period.

3.2 Performance on Pinheads and Sporophore Production

The effect of casing thickness on *Calocybe indica* production was also evaluated by measuring the number of pinheads and sporophores. The 1.0" casing thickness yielded the highest number of pinheads (28.60) and sporophores (12.6). In contrast, the 0.5" casing thickness resulted in the lowest counts, with 11.8 pinheads and 7.6 sporophores. The 0.75" and 1.5" casing thicknesses produced intermediate results, with 14.8 and 22.4 pinheads and 9.2 and 11.6 sporophores, respectively (Fig. 2). The limited number of pinheads and sporophores could be attributed to inadequate colonization and insufficient water retention capacity, highlighting the delicate balance needed for successful mushroom development.




Fig. 1. Effects of various levels of casing thickness on pin head initiation and sporophore maturation of *C.indica*

Table 1. Effect of casing material thickness on pinheads and fruiting body formation of *C.indica*

Casing Thickness	Days for pinhead formation	No. of pinheads	Days for fruiting body harvest	No. of fruiting body harvested
0.5"	17.40	11.80	14.40	7.60
0.75"	14.20	14.80	12.20	9.20
1.0"	10.40	28.60	9.00	12.60
1.5"	16.40	22.40	12.40	11.60
SEm (±)	0.48	1.18	0.64	0.48
CD @ 1%	1.44	3.55	1.93	1.44
CV %	7.35	13.64	12.00	10.46

Table 2. Effect of casing material thickness on stipe length, pileus, yield and biological efficiency of *C.indica*

Casing Thickness	Stipe length (cm)	Pileus diameter (cm)	Yield (g)	Biological efficiency (%)
0.5"	10.16	2.66	447.90	44.79
0.75"	9.76	2.90	591.50	59.15
1.0"	9.70	3.48	717.50	71.75
1.5"	8.92	3.22	674.10	67.41
SEm (±)	0.60	0.20	13.03	1.30
CD @ 1%	NS	NS	39.07	3.91
CV %	13.84	14.88	4.79	4.79




Fig. 2. Effects of different levels of casing thickness on pinhead initiation and sporophore harvest of *C.indica*

3.3 Performance on Yield and Biological Efficiency

The yield and biological efficiency of *Calocybe indica* varied significantly with different casing layer thicknesses, as detailed in Table 2. The 1.0" casing thickness produced the highest yield (717.5 g) and biological efficiency (71.75%),

followed by the 1.5" casing thickness, which resulted in a yield of 674.1 g and a biological efficiency of 67.41%. The 0.75" casing thickness yielded 591.5 g and a biological efficiency of 59.15%. Conversely, the 0.5" casing thickness resulted in the lowest yield (447.9 g) and biological efficiency (44.79%) (Fig. 3). Statistical analysis revealed no significant relationship

between stipe length and pileus diameter in relation to yield and biological efficiency (Table 2).

Casing is a critical step in milky mushroom production as it triggers the reproductive phase of the mushroom and facilitates substrate aeration. Among the various casing thicknesses tested, 1.0" proved to be the most effective, promoting early pinhead initiation, a shorter sporophore maturation period, higher yield, and greater biological efficiency under the red and lateritic belt conditions of West Bengal. This thickness also significantly outperformed other casing layers. These findings are consistent with those of Chiwan [9], who identified a casing thickness of 10-25 mm as ideal for prompt primordial initiation and sporophore maturation.

Although the 0.75" casing thickness also led to early pinhead initiation compared to the 1.5" thickness, it resulted in fewer pinheads and sporophores. The superior performance of the 1.0" thickness is attributed to its ability to provide adequate substrate aeration and facilitate the easy upward growth of pinheads. Similar conclusions were reached by Sharma et al. [10], Prasuna [11], and Shukla [12], all of whom observed that a 1.0" casing thickness was optimal for yield and biological efficiency in *Calocybe indica*. However, this result partially

contrasts with Krishnamoorthy's [13] findings, which suggested that a casing thickness less than 1.0" was preferable for achieving better yields of *C. indica*.

3.4 Effect of Casing Material

Casing, which involves covering the mycelial substrate with a layer of soil, is essential for supporting the mushroom canopy and ensuring proper fruiting. It signals the end of the vegetative growth phase and the commencement of mushroom reproduction. According to Phutela et al. [14], ideal casing material should be free from pests, diseases, competitor organisms, undecomposed vegetable matter, and should be neutral in pH.

The thickness of the casing material significantly influences primordial initiation, number of primordia, and overall yield. Krishnamoorthy [13] recommended a casing thickness of 1-2 cm for optimal yield. Conversely, Sharma et al. [10] and panda and Biswas [8] found that a 1.0" casing thickness was superior for the yield of *C. indica*. Prasuna [8] also supported the use of a 1.0" casing thickness for achieving the highest yield, compared to 1.5" and 2.0" thicknesses. Shukla [12] tested various casing thicknesses (0.5, 1.0, 1.5, 2.0, and 2.5 inches) and identified 1.0-1.5" as the most effective for maximum yield.




Fig. 3. Effects of various levels of casing material thickness on the yield and biological efficiency of *C.indica*

Pani [15] investigated different casing depths and timings and found that a thickness of 10-25 mm was ideal for primordia initiation and sporophore production. Chiwan [9] also reported that a casing thickness of 15-25 mm was optimal for rapid primordial initiation and sporophore maturation.

4. CONCLUSION

The introduction of *Calocybe indica* cultivation in West Bengal, offers a promising new avenue for agricultural expansion and farmer prosperity. This study revealed that a casing thickness of 1.0” is ideal for cultivating milky mushrooms, significantly enhancing primordia initiation, sporophore maturation, yield, and biological efficiency. By adopting this optimal casing thickness, local farmers can achieve higher yields and improved crop quality, potentially transforming their agricultural practices and boosting their incomes. These findings not only provide valuable guidance for commercial mushroom growers but also support the broader goal of diversifying agriculture in West Bengal, fostering economic growth, and improving the livelihoods of small and marginal farmers. The successful commercialization of milky mushrooms could thus play a crucial role in advancing rural development and sustainability in the region.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENTS

We would like to express our gratitude to the Department of Plant Pathology at Palli Siksha Bhavana, Visva-Bharati, for providing the facilities necessary to conduct this experiment. We also extend our thanks to everyone who contributed directly or indirectly to the success of this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Purkayastha PR, Chandra A. New species of edible mushroom from India. Trans Brit Mycol Societ. 1974;62:415-418.

2. Krishnamoorthy AS, Muthusamy M, Marimuthu T, Narasimhan V, Muthusankaranarayanan A. APK2 milky mushroom-Extn. Bulletin. RRS, TNAU, Aruppukottai; 1998.
3. Subbiah AK, Balan V. A Comprehensive Review of Tropical Milky White Mushroom (*Calocybe indica* P&C). Mycobiol. 2015;43(3):184–194. DOI: 10.5941/MYCO.2015.43.3.184
4. Ayimbila F, Keawsompong S. Nutritional Quality and Biological Application of Mushroom Protein as a Novel Protein Alternative. Current Nutrition Reports, 2023;12:281–298. DOI: 10.1007/s13668-023-00468-x
5. Ghosh S, Acharya K. Milky mushroom: A healthy nutritious diet. Food Res. Int. 2022; 156:111113. Available:https://doi.org/10.1016/j.foodres.2022.111113
6. Kumar S, Sharma VP, Shirur M, Kamal S. Status of milky mushroom (*Calocybe indica*) in India-a review. Mushroom Res. 2017;26(1):21-39.
7. Biswas MK. Oyster Mushroom Cultivation: a Women Friendly Profession for the Development of Rural West Bengal. Int J Bio-Res Stres Mangt. 2014;5(3):432-435. DOI:10.5958/0976-4038.2014.00594.6
8. Panda D and Biswas MK. Impact of sterilization and thickness of casing Materials on Yield attributes of *Calocybe indica*. Res. J Agrl. Sci. 2021;12(3):1004-1011.
9. Chiwan AA. Studies on the cultivation and productive potential of milky mushroom (*Calocybe indica* P&C) on different lignocellulosic wastes. Ph.D Thesis, Department of Botany. University of Jammu; 2014. Available:Http://Hdl.Handle.Net/10603/152614
10. Sharma SS, Doshi A, Trivedi A, Kothari KL. Studies on spawn research of *Calocybe indica* (P&C). Ind Bot Rep. 1997;14:97-100.
11. Prasuna K. Studies on nutritional and environmental factor effecting growth and yield of *Calocybe indica*. Osmania University. Hydrabad: Osmania University; 2002.
12. Shukla C. Role of Agronomical and Biochemical Parameters on Growth and Yield of *Calocybe indica* (P&C). Ph.D thesis, Department of Plant Pathology,

- Indhira Gandhi Krishi Visvavidhyalaya, Raipur, Chattisgarh, India; 2003.
13. Krishnamoorthy A. Studies on cultivation of milky mushroom (*C. indica*). Ph.D thesis, Tamil Nadu Agricultural University. Department of plant pathology, Coimbatore, Tamil Nadu, India; 1995.
14. Phutela UG, Phutela R. Effect of Physical and Chemical factors on growth of *Calocybe indica* (P & C). Int J Ad Life Sci. 2012;2:8-15.
15. Pani B. Sporophore production of milky mushroom (*Calocybe indica*) as influenced by depth and time of casing. Agrl Food Sci. 2012; 2, 168-170. Available: <https://api.semanticscholar.org/CorpusID:212562267>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124292>