



# Allelopathic Potential of Medicinal Plants Serves as a Valuable Source for Sustainable Weed Management

**Mohamed Salman M<sup>a++</sup>, Gurusamy A<sup>a##</sup>, Subramanian E<sup>bt</sup>,  
Prabhakaran J<sup>a‡</sup> and Arunachalam P<sup>c^</sup>**

<sup>a</sup> Department of Agronomy, Agricultural College & Research Institute, TNAU, Madurai, Tamil Nadu, 625104, India.

<sup>b</sup> ICAR-KVK, Agricultural College & Research Institute, TNAU, Madurai, Tamil Nadu, 625 104, India.

<sup>c</sup> Department of Plant breeding and genetics, Agricultural College & Research Institute, TNAU, Madurai, Tamil Nadu, 625 104, India.

## Authors' contributions

This work was carried out in collaboration among all authors. Author MSM conceptualized the review, outlined the key sections, and prepared the initial draft of the manuscript. Author GA and Author SE provided critical insights on specific topics, managed literature collection, and contributed to the writing of the thematic sections. Author PJ conducted a thorough review of the references, ensuring the inclusion of the most recent and relevant studies. Author AP reviewed and revised the manuscript for intellectual content and final approval. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijecc/2024/v14i104513>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124339>

**Review Article**

**Received: 20/07/2024**

**Accepted: 25/09/2024**

**Published: 10/10/2024**

<sup>++</sup> PG Scholar;

<sup>#</sup> Professor;

<sup>†</sup> Programme Coordinator;

<sup>‡</sup> Associate Professor (SS&AC);

<sup>^</sup> Associate Professor;

\*Corresponding author: E-mail: [gurusamy.a@tnau.ac.in](mailto:gurusamy.a@tnau.ac.in);

**Cite as:** Salman M, Mohamed, Gurusamy A, Subramanian E, Prabhakaran J, and Arunachalam P. 2024. "Allelopathic Potential of Medicinal Plants Serves As a Valuable Source for Sustainable Weed Management". *International Journal of Environment and Climate Change* 14 (10):633-48. <https://doi.org/10.9734/ijecc/2024/v14i104513>.

## ABSTRACT

Weeds are noxious plants that cause major problems for crop cultivation by interfering with crop growth and development. Due to weed infestation, about 50% of crop yields are reduced. In fact, weed causes more damage than pests and diseases in crop production. In modern agriculture, weeds are managed by using synthetic chemical herbicides. However, in organic farming, weed management is only possible through conventional methods such as manual and mechanical weeding, which are uneconomical and labor-intensive processes. Nowadays, labor shortage is the major problem in agriculture for performing various intercultural operations, especially weed management. In addition, continuous usage of synthetic herbicides can cause contamination to the surroundings thereby resulting in environmental pollution and the evolution of several herbicide-resistant weed species. To address these drawbacks allelopathic potential and allelochemicals can be utilized for weed management. Medicinal plants are rich sources of secondary metabolites with allelopathic potential, which helps to manage weeds by using plants extracts and natural herbicides. Allelochemicals of medicinal plants have the greatest potential for biological weed management by utilizing plant extracts and extracted allelochemicals as natural herbicides. Naturally extracted allelochemicals have almost the same effect as that of using synthetic herbicides in weed management without causing hazards to the environment. Thus, natural herbicides play a vital role in maintaining eco-friendly agriculture and sustainability in soil health. This review will help to create a general view about the allelopathic potential of medicinal plants and also helps to identify future research areas for weed management in organic farming with the help of allelochemicals.

*Keywords: Allelopathic potential; medicinal plants; natural herbicides; secondary metabolites; weeds.*

## 1. INTRODUCTION

Weeds are considered as the most important deleterious pest and they rank first in causing yield loss ranging from 33-55% in the crop production [1], [2], [3]. In a global context, this results in significant economic losses estimated at around USD 32 billion annually, primarily due to the impact of weeds [4]. In organic farming, farmers are facing major problems with economical and efficient weed management practices [5]. Organic farmers generally follow conventional methods viz. manual and mechanical weed management practices. But the major drawbacks of conventional weed management practices are uneconomical, labor-intensive, and time-consuming processes [6]. Further, in this decade, wage rates have drastically increased, and the availability of labor has drastically reduced [7], [8]. In modern agriculture, synthetic chemical herbicides are utilized for effective weed management: in fact, farmers use tons of synthetic chemical herbicides for weed management [9]. Moreover, the uncontrolled use of chemical herbicides causes environmental pollution and harmful effects on soil, water, animals, and humans [10], [11], [12]. The prolonged usage of chemical herbicides results in the emergence of new herbicide-resistant weed species. In fact, persistence in herbicide usage paves the way for the development of 310 weed species as herbicide-

resistant biotypes [13]. By adopting sustainable agriculture principles, the usage of chemical pesticides and herbicides can be reduced [14]. Recently, researchers have focused on the development of eco-friendly, natural herbicides [15], [16]. These natural herbicides are developed from microorganisms [17], [18] and plant-originated allelochemicals [19], [20]. Allelochemicals are known for causing negative or positive interactions on another organism through allelopathy [21]. The allelopathic potential of medicinal plants is effective in weed management without causing environmental issues [22], [23]. The use of plant-extracted allelochemicals in weed management helps to overcome labor shortages and environmental pollution [24]. According to Ray *et al.*, 2022 [8], allelopathic potential and allelochemicals are innovative solutions for sustainable weed management practices. This review will discuss the various importance and properties of allelochemical compounds in medicinal plants.

## 2. ALLELOPATHY AND ALLELOCHEMICALS

The harmful or beneficial effects of one plant on another due to the production of allelochemicals are referred to as "allelopathy" [25], [26]. It is also defined as a plant-to-plant chemical interaction that may either promote or interfere with the growth and development of plants present in the

surroundings [27]. Allelopathy effects are caused by some chemical compounds, particularly secondary metabolites, which are known as allelochemicals. These chemicals can produce both positive and negative interactions in the surrounding environment [28]. Generally, allelochemicals are non-nutritive and naturally originate from living organisms, especially microorganisms, plants, and animals [29].

Yankova-Tsvetkova *et al.*, [30] identified that the allelochemical compounds mainly consist of secondary metabolites such as alkaloids, coumarins, fatty acids, flavonoids, ketones, phenolics, purines, quinine, terpenoids, and aldehydes. Further, these allelochemicals are divided into 14 different groups based on their similarities. These 14 groups include aliphatic aldehydes, alkaloids, amino acids and peptides, anthraquinone, benzoic acid and its derivatives, benzoquinone, cinnamic acids and its derivatives, complex quinones, coumarins, cyanohydrins, flavonoids, glucosinolates, ketones, long-chain fatty acids, nucleosides, polyacetylenes, purines, simple phenols, steroids, terpenoids, and water-soluble organic acids [31]. For example, in *Eucalyptus* species, 18-cineole is a volatile terpene compound released as an allelochemical that inhibits the germination and growth of the weed *Amaranthus retroflexus* L. [32].

The plant parts such as leaves, fruits, seeds, flowers, stems, bark, and roots are responsible for the release of allelochemicals at different concentrations into the environment [33], [34], [35]. and differ in their pathway of interaction and release into the environment [36]. The following ways are responsible release of allelochemicals 1) through root exudation of chemical compounds 2) the release of allelochemicals from the decomposition of plant tissues and residues 4) releasing of volatile allelochemicals from living tissues of plants [21].

### 3. SCOPE FOR UTILIZATION OF ALLELOCHEMICALS IN WEED MANAGEMENT


There are several methods for utilization of allelochemicals in weed management practices such as intercropping, cover crops, mulching, crop rotation, plant extracts and natural herbicides as shown in Fig.1 [37]. Intercropping with allelopathic potential species can prevent

weed growth and the development by releasing allelochemicals into the soil [38], [39]. According to Saady and Science (2015) [40] weed growth has been reduced while using cowpea as an intercrop with maize due to competition and soil-released allelochemicals.

Crop rotation can significantly suppress the weed growth and disease-causing pathogens and pest incidence. Allelochemicals released from the crop rotation by roots and decomposition of crop residues helps in controlling weed seed germination [41]. Jabran and Chauhan. (2018) [42] recorded that weed infestation was reduced in the wheat crop, when the wheat crop was grown following the sorghum crop as a result of allelochemicals released from sorghum. Cover crops are also utilized in sustainable weed management as they possess allelopathic effects and causes competition for light, nutrients, water, and sunlight for weed growth. This result suppresses the weed population [39]. An allelochemical L-DOPA (L-3, 4-dihydroxyphenylalanine) was produced and identified in the field when velvet beans were grown as a cover crop [43]. Mulching with the residue of allelopathic potential plants is an effective source for sustainable weed management [44]. Using crop residue as mulches can suppress weed growth through allelochemicals released from crop residue [45], [46], [47]. According to Molina *et al.* (1991) [48] releasing of allelochemicals from Eucalyptus litter restrict the growth of many weeds. Most of the allelochemicals are secondary metabolites these compounds can be water soluble and water act as a medium of extraction for most of allelochemicals [37]. The Aqueous extracts of *Datura* consist of allelochemicals Scopolamine, hyoscyamine and atropine has the ability to inhibit the growth of *Parthenium hysterophorus* [49], [50]. In recent days the allelopathic potential of plants extracts can be utilized for the discovery of new natural herbicides [51]. A list of commercially developed natural herbicides with their natural allelopathic components of medicinal plants are given in "Table 1". This natural herbicide can reduce the usage of synthetic herbicides [52], [53]. There are two ways of extracting allelochemicals from plant tissue 1) identification and isolation of active chemical compounds and 2) Aqueous extract or organic extracts can be used directly for preparation of natural herbicides [54].

**Table 1. List of commercial developed natural herbicides and their responsible allelopathic components of medicinal plants [55], [56], [57]**

Allelopathic components of medicinal plants	Natural herbicide products
Citrus oil (70%)	Weed Zap™
Citronella grass oil (50%) and Wintergreen oil and clove oil (46%)	GreenMatch EX™
Cinnamon oil (30%)	Matran II™
Chinese perfume plant ( <i>Aglaia odorata</i> ) leaf extracts	Weed Zap™
	PORGANIC™



**Fig. 1. Methods of using allelochemicals in weed management**

#### 4. ALLELOPATHIC POTENTIAL OF MEDICINAL PLANTS AND HERBS IN WEED MANAGEMENT

In the world about 52,885 flowering plants are identified and evaluated as medicinal plants [58]. Generally, the chemical compounds produced from medicinal plants are used as healing agents and preparation of tablets, tonics, syrups [59], [60], [61]. From the total medicinal plants, only 6 percent of plants are examined for their biological and phytochemical properties [62]. From the examined medicinal plants about 40000 secondary metabolites were identified but only some secondary metabolites have been selected for their effective herbicidal activities [63]. Some medicinal plants with their phytotoxic effects against weeds are listed in “Table 2”.

Medicinal plant consists of large amounts of secondary metabolites; these secondary metabolites are responsible for performing various biological activities such as plant defense

against biotic and abiotic stress and the production of allelopathic potential compounds [71], [57], [72]. According to Islam and Kato-Noguchi. (2014) [73], the phytotoxic potential can be easily identified in medicinal plants which contain numerous bioactive and secondary metabolites than any other plant groups. Fujii and Appiah. (2018) [74] started detailed studies on the allelopathic potential of medicinal plants and identified that the screened compounds possess growth inhibition of some weed species. A list of identified medicinal allelopathic plants with their potential allelochemicals against the growth and development of weeds are given in “Table 3”. Azizi *et al.* (2009) [75] analyzed the allelopathic potential of 56 aromatic medicinal plants species in Iran and the study exposed that the 51 aromatic medicinal plants species possess inhibitory activity against the seedling growth of lettuce. The allelopathic potential of aromatic medicinal such as *Ocimum* spp, *Mentha* spp, *Leucas* spp, *Datura* spp and *Eucalyptus* spp are elaborately discussed.

**Table 2. List of medicinal plants with their phytotoxic effects against weeds**

Common name	Scientific name	Plant Parts	Targeted weed species	Effects on growth and the development	References
Common barberry	<i>Berberis vulgaris</i>	Whole plant extracts	Lettuce weed	80 percent suppression in growth	[64]
Saffron crocus	<i>Crocus sativus</i>	stigma and style	Lettuce weed	Strong inhibitory effects against seedlings	[65]
Golden apple	<i>Aegle marmelos</i>	Whole plant extracts	<i>Raphanus sativus</i>	100 percent inhibits the growth	[66]
Camelthorn	<i>Alhagi maurorum</i>	Aqueous extract	<i>Amaranthus retroflexus</i> , <i>Chenopodium murale</i>	Prevents the germination and growth of seedlings in laboratory experiment	[67]
Black Jack	<i>Bidens Pilosa</i>	Litter and residue incorporation @2tons ha <sup>-1</sup>	Rice ecosystem weeds	Reduce weeds growth up to 80% in rice and increase yield up to 20 percent.	[68]
Chinese Perfume Plant	<i>Aglaia odorata</i>	Leaves powder application @ 1ton ha <sup>-1</sup>	<i>Digitaria adscendens</i> , <i>Trianthema portulacastrum</i> and <i>Amaranthus gracilis</i>	Inhibited the emergence and growth	[69]
White Weed Black Jack Milk peas	<i>Ageratum conyzoides</i> , <i>Bidens pilos</i> , <i>Galactia pendula</i> .	Residue incorporation @ 1-2 tons ha <sup>-1</sup>	<i>Rotala indica</i>	Inhibits the emergence	[70]

**Table 3. List of identified medicinal allelopathic plants with their potential allelochemicals against growth and development of weeds**

Common name	Scientific name	Family	Allelochemical Compound isolated	Phytotoxicity effects on targeted plants	Targeted weed species plants	References
Pignut	<i>Hyptis suaveolens</i>	Lamiaceae	suaveolic acid (14 $\alpha$ -hydroxy-13 $\beta$ -abiet-8-en-18-oic acid)	Inhibition of growth and development	Italian grass, barnyard grass, garden cress	[76]
Neem	<i>Azadirachta indica</i>	Meliaceae	Nimbolide B and Nimbic acid B.	Root and shoot growth reduction	Barnyard grass ( <i>Echinochloa</i> spp)	[77]
Neem	<i>Azadirachta indica</i>	Meliaceae	Nimbolide B and Nimbic acid B.	Inhibits germination	<i>Lepidium sativum</i> , <i>Latuca sativa</i>	[78]
Garden asparagus	<i>Asparagus officinalis</i>	Asparagaceae	trans-Cinnamic acid, p-Coumaric acid and iso-Agatharesinol	Inhibits the root and shoot growth	Rye grass	[79]
Prickly ash	<i>Zanthoxylum limonella</i>	Rutaceae	Brevifolin	Inhibits the seed germination and growth	Chinese amaranthus	[80]
<b>Thorn-Apple</b>	<i>Datura metel</i>	Solanaceae	hyoscyamine, atropine	Reduce emergence and growth	<i>Parthenium</i> spp	[49]
River Red Gum tree	<i>Eucalyptus camaldulensis</i>	Myrtaceae	p-coumaric, gallic, gentisic,p-hydroxybenzoic, syringic acid, catechol and vanillic acid,	Suppression of germination and growth	<i>Portulaca oleracea</i>	[81], [82]

Common name	Scientific name	Family	Allelochemical Compound isolated	Phytotoxicity effects on targeted plants	Targeted weed species plants	References
Bluegum eucalyptus	<i>Eucalyptus globulus</i>	Myrtaceae	Hyperoside, kaempferol 3-O-glucoside, shikimic-succinic acids	Inhibit germination and physiological growth.	<i>Agrostis stolonifera</i>	[83]
Long Pepper	<i>Piper longum</i>	Piperaceae	Sarmentine	Inhibits growth	<i>Convolvulus arvensis</i> <i>Conyza canadensis</i> <i>Sinapis arvensis</i>	[84]
Aleo vera	<i>Aloe barbadensis</i>	Liliaceae	Anthraquinones	Inhibits the germination and seedling growth	Dandelion <i>Taraxicum officinalis</i> )	[85]

**Ocimum spp.:** Generally, the *Ocimum* genus has been classified into 65 different species [86] and is widely distributed in tropical and subtropical of Asia [87], [88]. Most popular species such as *Ocimum tenuiflorum* L., *Ocimum sanctum* L., *Ocimum gratissimum* L., *Ocimum americanum* L. (*O. canum* S.), *Ocimum basilicum* L., *Ocimum kilimandscharicum* G. and *Ocimum micranthum* W. are grown for their incredible medicinal properties. In *ocimum* majority of allelopathic bioactive compounds are grouped as terpenoids in the composition of Terpene alcohol (linalool); Acyclic monoterpenes (alloocimene, p-cymene, citronellal, cis- $\beta$ -ocimene); Sesquiterpenes ( $\beta$ -carophyllene); Monoterpene bicyclic ether (1,8-cineole); Cyclic monoterpenes (limonene); Bicyclic monoterpenes (camphene,  $\alpha$ -pinene); Terpenic hydrocarbons, and fewer diterpenes [89]. *Ocimum basilicum* L. commonly known as sweet basil is the most vital aromatic herbs composed of a variety of secondary metabolites such as Flavonoids, alkaloids, terpenoids, essential oil, tannin, phenols, steroids, saponins, aldehydes, and glycosides [90]. Basil (*O. sanctum* L) is a widely grown medicinal plant also Grown for their religious belief in India. It is classified under the Lamiaceae family; many secondary metabolites such as steroid ursolic acid and n-triacontanol are extracted from the basil leaves. The potential allelopathic compounds such as Methyl ether (4.8%), Eugenol (70.5%), Apinene (3.5%), Caryophyllene (7.5%), Terpinen-4-(0.4%), Nerol (6.4%), Terpinen-4-(0.4%), Selenine (0.4%), Pinene (0.4%), and Decyl aldehyde [91], [89]. Hiltunen and holm. [92] identified that the *O. tenuiflorum* plant extracts consist of phytotoxic substances such as methyl chavicol, linalool, methyl eugenol and methyl cinnamate in addition to some subtypes of oils. Kamel et al. [93] identified some allelopathic bioactive compounds with GC-MS analysis in *Ocimum Sp*, which contains m-Camphorene (5.45%), Phenol, 2-methoxy-3-(2-propenyl)- (9.79%), Ethyl-3-hydroxyandrostane-17-one (4.62%), Geranyl- $\alpha$ -terpinene (6.87%), Retinol, acetate (6.77%), 1,3,6,10-Cyclotetradecatetraene, 14-Isopropyl-3,7,11-Trimethyl(+)- (8.69%), Eugenol (9.79%), 2-(7-Heptadecyloxy)tetrahydro-2H-pyran (4.62%), 1,3,6,10 Cyclotetradecatetraene, 14-Isopropyl-3,7,11-Trimethyl(+)- and p-Camphorene (5.45%) are the compounds identified. Singh et al. [94] observed that the *Parthenium hysterophorus* seedling growth is inhibited by plant extracts of *Ocimum americanum*. The 30 percent aqueous concentration extract of *Ocimum sanctum* has

shown 70% inhibition of seed germination of *Parthenium hysterophorus* [95]. According to Dhima et al. [96] *Ocimum basilicum* incorporation in the soil as green manure inhibits the emergence of *Echinochloa sp* (barnyard grass) by 11-50 percent. The reduction in the total weed population in cowpea and groundnut has been observed by the spraying of plant extracts of *Ocimum basilicum* L as pre-emergence [97]. The aqueous extract of *Ocimum basilicum* has the potential to control weeds such as *Amaranthus*, jungle rice, galium, and common purslane, and also supports the growth of groundnut and cowpea. Moreover, *Ocimum basilicum* extracts may be used as natural herbicides for sustainable weed management [91]. *Ocimum americanum* contains allelochemicals such as limonene, camphor and linalool have the potential to restrict the germination and seedling growth weed *Mimosa pudica* and *Senna obtusifolia* [98].

**Mentha spp.:** The *Mentha* genus is widely distributed in Northeastern Africa, western Asia and southeastern Europe [99]. *Mentha* genus belongs to the Lamiaceae family with 13 different species and the most common species with allelopathic activities are *Mentha spicata*, *Mentha suaveolens* and *Mentha piperita* [100]. In *Mentha suaveolens* (apple mint) more than 32 secondary metabolites are identified from which active allelochemicals such as 16 monoterpenes (Camphene,  $\alpha$ -pinene,  $\beta$ -ocimene, (+)-Sabinene,  $\beta$ -myrcene,  $\beta$ -thujene, Tricyclene, D-limonene,  $\gamma$ -terpinene,  $\alpha$ -terpinene, p-menthan-1-ol, Linalool, 4-thujanol, Endo-borneol and Terpinen-4-ol) and 5 Sesquiterpenes ( $\alpha$ -bourbonene, Caryophyllene, Cadina-1(6),4-diene,  $\beta$ -cubebene 2 and Germacrene) [101]. Mahdavia et al. [102] have founded that GC-MS analysis of *Mentha piperita* L. (peppermint) revealed that the presence of allelochemicals such as 8-cineole (5.9%), mentone (17.48%), menthol (35%), and menthofuran (11.7%). Additionally, the extracts also contain some determined phenolic compounds were found namely trans-ferulic acid (10.8 mg/g), hesperidin (9.3 mg/g), ellagic acid (6.8 mg/g), and sinapic acid (4.2 mg/g). According to Campiglia et al. [103] seed germination of ryegrass was significantly inhibited by peppermint oil. The seedling growth of barnyard grass and *Timothy sp* are inhibited by using the aqueous methanol extracts of *Mentha sylvestris* [26]. According to Petrova et al. [104]; Nagaraja and Deshmukh. [105] *Mentha spicata* oil exhibit the allelopathic potential against some weeds namely

*Amaranthus retroflexus*, *Sinapis arvensis*, *Centaurea salsotitalis*, *Sonchus oleraceus*, *Rumex nepalensis*, *Raphanus raphanistrum* and its extracts shows inhibitory effects on *Cynodon dactylon*, *Rumex crispus* and *Sorghum halepense*. The allelopathic effects of *Mentha piperita* extracts and its essential oil inhibit the seed germination (%), root and shoot growth and dry weight of weeds such as field bindweed (*Convolvulus arvensis* L.), purslane and jungle rice (*Echinochloa colonum* L.) [106], [107]. The seed germination of *Amaranthus* spp. and *E. colona* are restricted by using essential oil produced from *Mentha spicata* (spearmint) [108]

**Leucas spp.:** The genus *Leucas* is classified into 80 different species of which more than 41 species are available in India [109]. The two major species *Leucas aspera* and *Leucas cephalotes* are widely distributed in the world. *Leucas aspera* is a popular aromatic medicinal herb that belongs to the family Lamiaceae and consists of numerous bioactive compounds. It is a common weed species found in fallow land and roadsides widely distributed in both tropical and temperate regions of Asia [110], [111]. Secondary metabolites such as terpenes, flavonoids, coumarins, steroids and lignans are present [112], [113], [111]. In particular, the methanolic extracts of *Leucas aspera* contains allelochemical compounds such as diterpenes (rel5S, 6R, 8R, 9R, 10S, 13S, 15S, 16R)- 6 acetoxyl-9,13;15, 16diepoxy-15hydroxy-16methoxyabdane and (rel5S, 6R, 8R, 9R, 10S, 13S, 15R,16R)- 6-acetoxyl-9,13;15,16-diepoxy-15-hydroxy methoxyabdane [76]. The growth and seed germination of *Echinochloa colona* was significantly inhibited by applying methanol extracts of *Leucas aspera* [114],[115]. According to Thang Lam and Hisashi. [116] methanol extracts of *L. cephalotes* exhibited inhibitory and allelopathic effects on barnyard grass, *Timothy sp*, Italian ryegrass.

**Datura spp.:** The genus *Datura* is a most popular medicinal plant used throughout the world and also it contains higher allelopathic effects against crops and weeds [49]. The two major species with allelopathic activities are *Datura metel* L and *Datura stramonium*. Ma et al. [117] have identified that the *Datura metel* contains phytotoxic compounds in the form of withanolide glycosides namely daturametelins H–J, daturaturin-A, and 7,27-dihydroxy-1-oxowitha-2,5,24-trienolide present in the aerial parts. Some other allelochemical compounds such as pyrrole compound derivative, 2-(3,4-

dimethyl-2,5-dihydro-1H-pyrrol-2-yl)-10 - methyl ethyl pentanoate was extracted and identified from the leaves of *Datura metel* [118] and sphingosine compound derivatives namely (4E,8Z)-1-O (-D-glucopyranosyl)-N-(20 hydroxyhexadecanoyl)-sphinga-4,8-dienine [119]. These compounds exhibit allelopathic herbicidal activity. The aqueous extracts of *Datura metel* have recorded inhibitory effects in germination and seedling growth of parthenium weed in laboratory assay, In the field evaluation the reduction in germination was observed up to 97 percent of parthenium is observed by incorporation of *D. metel* residue at the rate of 4- 5 percent [49].

*Datura stramonium* is a small herb of 1m in height and is widely distributed all over the world [120].

*D. stramonium* is another important species that contains numerous allelochemicals such as atropine, scopolamine and hyoscyamine [121]; and these allelochemicals inhibit the root and shoot growth of weeds namely *Trigonella* and *Lepidium* [122]. In *D. stramonium* 44 allelopathic bioactive compounds have been identified by using GC-MS analysis, compounds namely sterols and their related compounds 26,26-Dimethyl-5,24(28)-ergostadien-3. beta-ol,5. alpha. -Ergosta-7,22-dien-3. beta and 3-Hydroxycholestan-5-Acetate [123]. Lovett et al. [124] identified some alkaloids derivatives responsible for restriction of seed germination and seedling growth of weeds such as *Cenchrus ciliaris*, *Notonia wightii*. A new natural herbicide can be derived from methanol extracts of *D. stramonium*, these herbicides can be effectively used as both pre-emergence and post-emergence weed management herbicides against *Tagetes minuta* and *Amaranthus hybridus* in higher concentrations [125].

**Eucalyptus spp.:** The *Eucalyptus sp* belongs to the family Myrtaceae and is mostly cultivated for paper production in addition it contains many medicinal values and is distributed widely in the Mediterranean regions of the world [126], [127]. The two major species are *E. globulus Labill.* and *E. camaldulensis Dehnh* [128] are widely cultivated. May and Ash. [129] reported that allelochemicals such as phenolic acids and terpenes are available in leaves barks and roots of *Eucalyptus* spp. The leaves contain many phenolic and volatile compounds capable of producing allelopathic effects on weeds [130], [127]. The existence of several allelopathic



terpenes compounds namely limonene, 1,8-cineol,  $\alpha$ - and  $\beta$ -pinene has been identified from the foliage of *Eucalyptus* spp [131]. Putnam. (1984) [132] observed that compound such as benzoic, cinnamic and phenolic acids are released from *Eucalyptus* species and they are capable of suppressing the growth of some neighboring crops and weeds. Iqbal *et al.* [133] analyzed the essential oil of *E. camaldulensis* and reported that 16 allelopathic potential compounds were identified, the major constituents of compounds namely, 1-8 cineole,  $\alpha$ -pinene, p-cymene and  $\beta$ -phellandrene. Due to the application of leachates extracted from fresh leaves of blue gum Eucalyptus (*E. globulus* Labill.) at the rate of 20 percent has a significant effect in reducing the rate of resprouting of purple nutsedge (*Cyperus rotundus* L.) by 57-68 percent [134] and also when this leachates extracts applied at the rate of 25 percent the growth of bermudagrass is suppressed by 66 percent in the greenhouse experiment [135]. According to Azizi and Fuji [136] observed that 80 percent of germination of *Amaranthus retroflexus* L. and 90 percent *Portulaca oleracea* L. are reduced by the application of *E. globulus* essential oils applied at 0.2 percent. In addition, aqueous extract of *E. citriodora* inhibits the germination and growth of parthenium seeds. In *Echinochloa colona* the application of *E. globulus* dry leaves in soils can reduce the number of branches produced by 59 percent and also reduce the fresh weight of weed by 66 percent [137].

## 5. LIMITATIONS OF USING ALLELOCHEMICALS AND NATURAL HERBICIDES

Although the identification and selection of allelochemicals from medicinal plants are comparatively easier than compared to other groups of plants, but the extraction of allelochemicals and synthesis of natural herbicides are very expensive, laborious, difficult, and time-intensive processes [16]. For example, from the allelochemicals cyclic tetrapeptide tentoxin has ability to synthesis natural herbicide but the extraction and isolation methods are very expensive and tedious processes [138], [39] and some allelochemicals like aflatoxin and ricin exhibits toxic to living organisms especially human and animals [139]. Moreover, the shelf-life of active allelochemicals is less, than compared to synthetic chemicals [140], [2]. According to Maurya *et al.* (2022) [2], allelochemicals composition and quality varies even in the same locations and taxonomic

groups of plants. In addition, the major limitations in using allelopathic potential and allelochemicals are delay in the suppression of weeds in field conditions and the information about mode action in targeted plants by using allelochemicals and natural herbicides are not yet identified but, in contrast distinct mode of action using synthetic herbicides in weeds and targeted plants are studied and identified in detail [8]. The Final and foremost limitation on natural herbicides and allelopathic potential is in the laboratory and controlled conditions the allelochemicals contain greater phytotoxicity on targeted plants but in field experiments while using the same allelochemicals the phytotoxicity ability is very much decreased due to various physical, chemical, and biological factors of the environment [141], [142], [143].

## 6. FUTURE RESEARCHABLE AREAS

1. In the future more researches are needed for identification and isolation of the potential allelochemicals for weed management in the field conditions.
2. In addition to weed management the pest and disease-controlling ability of synthesized natural herbicides should be identified and studied to increase the efficacy in their application.
3. More studies are needed on physiology changes, mode of action, and bioassay of targeted plants by using allelochemicals.

## 7. CONCLUSION

From this review, it is clear that the medicinal plants are capable of producing several secondary metabolites with allelopathic potential and these secondary metabolites can be utilized as potent source for sustainable weed management as allelochemicals. The allelochemicals have the ability to reduce and suppress weed growth and development. These compounds after extraction and identification can be effectively used for development of new natural herbicide. The developed natural herbicides can be used as source for weed management in organic farming. It helps to maintain sustainability and pave the way for eco-friendly farming without affecting the environment.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that generative AI technologies such as Large Language Models,

etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. GRAMMARLY WAS USED TO CHECK GRAMMAR MISTAKES

## ACKNOWLEDGEMENT

The authors gratefully acknowledge the financial support provided by the Venture Capital Scheme of the Department of Agronomy, Agricultural College and Research Institute, Madurai, Tamil Nadu, India. We also extend our thanks to the DST-FIST (Department of Science and Technology - Fund for Improvement of S&T Infrastructure) scheme for providing infrastructural support, which was instrumental in the successful completion.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Siddiqui I, Bajwa R, Huma Z, Javaid A. Effect of six problematic weeds on growth and yield of wheat. *Pakistan J Bot.* 2010;42(4):2461-71.
2. Maurya P, Mazed A, Kumar D, Ahmad IZ, Suryavanshi P. Medicinal and aromatic plants as an emerging source of bioherbicides. *Curr Sci.* 2022;122(3):258-66. DOI:10.18520/cs/v122/i3/258-266.
3. Benjamin J, Idowu O, Babalola OK, Oziegbe EV, Oyedokun DO, Akinyemi AM, Adebayo A. Cereal production in Africa: the threat of certain pests and weeds in a changing climate—a review. *Agric Food Secur.* 2024;13(1):18.
4. Kubiak A, Wolna-Maruwka A, Niewiadomska A, Pilarska AA. The problem of weed infestation of agricultural plantations vs. the assumptions of the European biodiversity strategy. *Agronomy.* 2022;12(8):1808.
5. Soni JK, Choudhary VK, Singh PK, Hota S. Weed management in conservation agriculture, its issues and adoption: A review.
6. Hozayn M, Monem A, Lateef E. Crop residues, an effective tool for improving growth of wheat and suppression of some associated weeds. *Asian J Crop Sci.* 2011;27(2):237-44.
7. Duary B, Mukherjee A. Distribution pattern of predominant weeds in wet season and their management in West Bengal, India.
8. Ray P, Bhowmick MK, Ghosh RK, Kumar S. Herbal herbicide: A low-cost and eco-friendly tool for weed management in smallholder farming. *Indian J Weed Sci.* 2022;54(1):73-9. DOI:10.5958/0974-8164.2022.00073.9.
9. Mesnage R, Székács A, Zaller JG. Herbicides: Brief history, agricultural use, and potential alternatives for weed control. *Herbicides.* 2021;1-20.
10. Wilson C, Tisdell C. Why farmers continue to use pesticides despite environmental, health, and sustainability costs. *Ecol Econ.* 2001;39(3):449-62. DOI:10.1016/S0921-8009(01)00238-5.
11. Duary B. Recent advances in herbicide resistance in weeds and its management. *Indian J Weed Sci.* 2008;40(3&4):124-35.
12. Aktar W, Sengupta D, Chowdhury A. Impact of pesticide use in agriculture: Their benefits and hazards. *Interdiscip Toxicol.* 2009;2(1):1-12. DOI:10.2478/v10102-009-0001-7.
13. Sharma V, Ghosh T, Sahil K, Guleria A, Thakur A, Rana G, Thakur M. Biotechnological interventions for pesticide remediation and soil health reclamation. In: *Microbial and Biotechnological Interventions in Bioremediation and Phytoremediation.* 2022;501-39.
14. Velten S, Leventon J, Jager N, Newig J. What is sustainable agriculture? A systematic review. *Sustainability.* 2015;7(6):7833-65. DOI:10.3390/su7067833.
15. Akbar M, Javaid A, Ahmad E, Javed T, Clary J. Holadysenterine, a natural herbicidal constituent from *Drechslera australiensis* for management of *Rumex dentatus*. *J Agric Food Chem.* 2014;62(2):368-72. DOI:10.1021/jf403565g.
16. Islam AM, Yeasmin SS, Qasem JR, Juraimi AS, Anwar MP. Allelopathy of medicinal plants: Current status and future prospects in weed management. *Adv Sci.* 2018. DOI:10.4236/as.2018.912110.

17. Bashir U, Khan A, Javaid A. Herbicidal activity of *Aspergillus niger* metabolites against parthenium weed. *Planta Daninha*. 2018;36.
18. Javaid A, Jubeen T, Khan IH, Jabeen K, Akbar M. Herbicidal potential of *Alternaria citri* Ellis and Pierce metabolites against *Parthenium hysterophorus* L. *Allelopathy J*. 2022;55(1):25-34.
19. Akbar M, Javaid A, Khalil T, Iqbal MS. Isolation of herbicidal compounds from *Melia azedarach* L. to control *Rumex dentatus* L. in wheat. *Allelopathy J*. 2022;55(2):163-76.
20. Erida G, Ichsan CN, Kurniawan T, Khan IH, Javaid A. Potential of secondary metabolites of *Ageratum conyzoides* L. in weed management: A review. *Allelopathy J*. 2023;58(1).
21. Chon S, Jennings J, Nelson CJ. Alfalfa (*Medicago sativa* L.) autotoxicity: Current status. *Allelopathy J*. 2006;18(1):57.
22. Hong NH, Xuan TD, Eiji T, Hiroyuki T, Mitsuhiro M, Khanh TD. Screening for allelopathic potential of higher plants from Southeast Asia. *Crop Protect*. 2003;22(6):829-36.
23. Javaid N, Shah MH, Khan IH, Javaid A, Waleed SM. Herbicidal activity of *Ageratum conyzoides* against parthenium. *Pakistan J Weed Sci Res*. 2020;26(2):137-46.
24. Muhammad Z, Inayat N, Majeed A, Ali H, Ullah K. Allelopathy and Agricultural Sustainability: Implication in weed management and crop protection—An overview. *Eur J Ecol*. 2019;5(2):54-61. DOI:10.2478/eje-2019-0014.
25. Molisch H. Einfluss einer pflanze auf die andere, allelopathie; 1937.
26. Islam AM, H. *Mentha sylvestris*: A potential allelopathic medicinal plant; 2013.
27. Fernandez C, Monnier Y, Santonja M, Gallet C, Weston LA, Prévosto B, Bousquet-Mélou A. The impact of competition and allelopathy on the trade-off between plant defense and growth in two contrasting tree species. *Front Plant Sci*. 2016;7:186358. DOI:10.3389/fpls.2016.00594.
28. Del Fabbro C, Gusewell S, Prati D. Allelopathic effects of three plant invaders on germination of native species: A field study. *Biol Invasions*. 2014;16:1035-42. DOI:10.1007/s10530-013-0555-3.
29. Lim CJ, Basri M, Ee GCL, Omar D. Phytoinhibitory activities and extraction optimization of potent invasive plants as eco-friendly weed suppressant against *Echinochloa colona* (L.) Link. *Ind Crops Prod*. 2017;100:19-34. DOI:10.1016/j.indcrop.2017.01.025.
30. Yankova-Tsvetkova E, Nikolova M, Aneva I, Stefanova T, Berkov S. Germination inhibition bioassay of extracts and essential oils from plant species. *Comptes rendus de l'Académie bulgare des Sciences*. 2020;73(9):1254-9.
31. Cheng F, Cheng Z. Research progress on the use of plant allelopathy in agriculture and the physiological and ecological mechanisms of allelopathy. *Front Plant Sci*. 2015;6:160714. DOI:10.3389/fpls.2015.01020.
32. Azizi M, Fuji Y. Allelopathic effect of some medicinal plant substances on seed germination of *Amaranthus retroflexus* and *Portulaca oleraceae*. *Acta Hort*. 2005;699:59-66. doi:10.17660/ActaHortic.2006.699.5.
33. Bertin C, Yang X, Weston LA. The role of root exudates and allelochemicals in the rhizosphere. *Plant Soil*. 2003;256(1):67-83. DOI:10.1023/A:1026290508166.
34. Tesio F, Ferrero A. Allelopathy, a chance for sustainable weed management. *Int J Sustainable Dev Ecol*. 2010;17(5):377-89. DOI:10.1080/13504509.2010.507402.
35. Motmainna M, Juraimi AS, Ahmad-Hamdani MS, Hasan M, Yeasmin S, Anwar MP, Islam AM. Allelopathic potential of tropical plants—a review. *Agronomy*. 2023;13(8):2063. DOI:10.3390/agronomy13082063.
36. Sathishkumar A, Srinivasan G, Ragavan T, Thiyageshwari S, Aananthi N. Allelopathic effect of different intercropping systems and tree leaf extract spray on weed density, dry matter, and weed control efficiency in irrigated cotton. *Int J Current Microbiol App Sci*. 2017;6(6):1322-9. DOI:10.20546/ijcmas.2017.606.155.
37. Farooq M, Khaliq A, Cheema Z, Cheema S. Application of allelopathy in crop production: Success story from Pakistan. In: Narwal SS, Zeng X, editors. *Allelopathy: Current trends and future applications*. Springer; 2013. p. 113-43. DOI:10.1007/978-3-642-30595-5\_6.
38. Baumann DT, Bastiaans L, Kropff MJ. Intercropping system optimization for yield, quality, and weed suppression combining

- mechanistic and descriptive models. *Agron J*. 2002;94(4):734-42.  
DOI:10.2134/agronj2002.7340.
39. Khamare Y, Chen J, Marble SC. Allelopathy and its application as a weed management tool: A review. *Front Plant Sci*. 2022;13:1034649.  
DOI:10.3389/fpls.2022.1034649.
  40. Saudy HS. Maize–cowpea intercropping as an ecological approach for nitrogen-use rationalization and weed suppression. *Arch Agron Soil Sci*. 2015;61(1):1-14.  
DOI:10.1080/03650340.2014.920499.
  41. Peters RD, Sturz AV, Carter MR, Sanderson JB. Developing disease-suppressive soils through crop rotation and tillage management practices. *Soil Tillage Res*. 2003;72(2):181-92.  
DOI:10.1016/S0167-1987(03)00087-4.
  42. Jabran K, Chauhan B. Non-chemical weed control. Academic Press; 2018.
  43. Fujii Y, Shibuya T, Yasuda T. Allelopathy of velvetbean: Its discrimination and identification of L-DOPA as a candidate of allelopathic substances. *Japan Agric Res Q*. 1992;25(4):238-47.
  44. Tiquia SM, Lloyd J, Herms DA, Hoitink HA, Michel Jr FC. Effects of mulching and fertilization on soil nutrients, microbial activity, and rhizosphere bacterial community structure determined by analysis of TRFLPs of PCR-amplified 16S rRNA genes. *Appl Soil Ecol*. 2002;21(1):31-48.  
DOI:10.1016/S0929-1393(02)00040-9.
  45. White RH, Worsham AD, Blum U. Allelopathic potential of legume debris and aqueous extracts. *Weed Sci*. 1989;37(5):674-9.  
DOI:10.1017/S0043174500072623.
  46. Batish DR, Singh H, Kaur S. *Crop allelopathy* and its role in ecological agriculture. *J Crop Prod*. 2001;4(2):121-61.  
DOI:10.1300/J144v04n02\_03.
  47. Shafique S, Javaid A, Shafique S. Management of littleseed canarygrass (*Phalaris minor* Retz.) by extracts and dry leaf biomass of *Parthenium hysterophorus* L. *Philippine Agric Sci*. 2013;96(4):426-31.
  48. Molina A, Reigosa M, Carballeira A. Release of allelochemical agents from litter, throughfall, and topsoil in plantations of *Eucalyptus globulus* Labill in Spain. *J Chem Ecol*. 1991;17:147-60.  
DOI:10.1007/bf00994428.
  49. Javaid A, Shafique S, Shafique S. Herbicidal effects of extracts and residue incorporation of *Datura metel* against parthenium weed. *Nat Prod Res*. 2010;24(15):1426-37.  
DOI:10.1080/14786410903075440.
  50. Elisante F, Ndakidemi P. Allelopathic effect of *Datura stramonium* on the survival of grass and legume species in the conservation areas. *Asian J Res Crop Sci*. 2014;2(1):27-43.
  51. Nornasuha Y, Ismail B. Sustainable weed management using allelopathic approach. *Malaysian Agricultural Bulletin*. 2017; 46(2).
  52. Motmainna M, Juraimi A, Uddin MK, Asib NB, Islam A, Hasan M. Assessment of allelopathic compounds to develop new natural herbicides: A review. *Allelopathy Journal*. 2021;52:21-40.  
DOI:10.26651/allelo.j/2021-52-1-1305.
  53. Motmainna M, Juraimi AS, Uddin MK, Asib NB, Islam AM, Hasan M. Bioherbicidal properties of *Parthenium hysterophorus*, *Cleome rutidosperma*, and *Borreria alata* extracts on selected crop and weed species. *Agronomy*. 2021;11(4):643.  
DOI:10.3390/agronomy11040643.
  54. Yang RZ, Tang CS. Plants used for pest control in China: A literature review. *Economic Botany*. 1988;42:376-406.
  55. Cornes D. Callisto: A very successful maize herbicide inspired by allelochemistry. *Weed Research*. 2005;45(6):356-359.
  56. Kato-Noguchi H, Suzuki M, Noguchi K, Ohno O, Suenaga K, Laosinwattana C. A potent phytotoxic substance in *Aglaia odorata* Lour. *Chemistry & Biodiversity*. 2016;13(5):549-554.
  57. Dayan FE, Cantrell CL, Duke SO. Natural products in crop protection. *Bioorganic & Medicinal Chemistry*. 2009;17(12):4022-4034.  
DOI:10.1016/j.bmc.2009.01.046.
  58. Wakdikar S. Global health care challenge: Indian experiences and new prescriptions. *Electronic Journal of Biotechnology*. 2004;7(3):02-03.
  59. Kamboj VP. Herbal medicine. *Current Science*. 2000;78(1):35-39.
  60. Verma S, Singh SP. Current and future status of herbal medicines. *Veterinary World*. 2008;1(11):347.
  61. Mukherjee PK, Ponnusankar S, Venkatesh M. Ethnomedicine in complementary.
  62. Appiah KS, Mardani HK, Osivand A, Kpabitey S, Amoatey CA, Oikawa Y, Fujii Y. Exploring alternative use of medicinal

- plants for sustainable weed management. *Sustainability*. 2017;9(8):1468. DOI:10.3390/su9081468.
63. Einhellig FA, Leather GR. Potentials for exploiting allelopathy to enhance crop production. *Journal of Chemical Ecology*. 1988;14:1829-1844. DOI:10.1007/BF01013480.
  64. Mardani H, Azizi M, Osivand A, Fujii Y. Evaluation of allelopathic activity of Iranian medicinal plants by sandwich method. *Journal of Weed Science and Technology*. 2014;53:85.
  65. Amini S, Azizi M, Joharchi MR, Moradinezhad F. Evaluation of allelopathic activity of 68 medicinal and wild plant species of Iran by the sandwich method. *International Journal of Horticultural Science and Technology*. 2016;3(2):243-253. DOI:10.22059/ijhst.2016.62923.
  66. Islam AM, Hasan M, Musha MMH, Uddin MK, Juraimi AS, Anwar MP. Exploring 55 tropical medicinal plant species available in Bangladesh for their possible allelopathic potentiality. *Annals of Agricultural Sciences*. 2018;63(1):99-107. DOI:10.1016/j.aosas.2018.05.005.
  67. Qasem J. Allelopathic effects of selected medicinal plants on *Amaranthus retroflexus* and *Chenopodium murale*. *Asian Journal of Plant Sciences*. 2002;10(2):105-122.
  68. Hong NH, Xuan TD, Eiji T, Khanh TD. Paddy weed control by higher plants from Southeast Asia. *Crop Protection*. 2004;23(3):255-261. DOI:10.1016/S0261-2194(03)00051-6.
  69. Laosinwattana C, Teerarak M, Charoenying P. Effects of *Aglaia odorata* granules on the seedling growth of major maize weeds and the influence of soil type on the granule residue's efficacy. *Weed Biology and Management*. 2012;12(3):117-122. DOI:10.1111/j.1445-6664.2012.00444.x.
  70. Xuan TD, Shinkichi T, Khanh TD, Chung IM. Biological control of weeds and plant pathogens in paddy rice by exploiting plant allelopathy: An overview. *Crop Protection*. 2005;24(3):197-206. doi:10.1016/j.cropro.2004.08.004.
  71. Wink M. Introduction: Biochemistry, role, and biotechnology of secondary metabolites (From APR Volume 2). *Annual Plant Reviews*. 2018:1-15. DOI:10.1002/9781119312994.apr0012.
  72. Rice E. *Allelopathy*. Orlando, FL: Academic Press; 1984.
  73. Islam A, Kato-Noguchi H. Phytotoxic activity of *Ocimum tenuiflorum* extracts on germination and seedling growth of different plant species. *The Scientific World Journal*. 2014. DOI:10.1155/2014/676242.
  74. Fujii Y, Appiah KS. Allelopathy for sustainable weed management. In: Labrada K, Gillespie JJ, editors. *Weed control*. CRC Press; 2018. p. 166-190.
  75. Azizi M, Amini S, Joharchi MR, Oroojalian F, Baghestani Z. Genetic resources for allelopathic and medicinal plants from traditional Persian experience. In: MARCO Symposium: Challenges for Agro-Environmental Research in Monsoon Asia; 2009. p. 38-46.
  76. Islam AM, Ohno O, Suenaga K, Kato-Noguchi H. Two novel phytotoxic substances from *Leucas aspera*. *Journal of Plant Physiology*. 2014;171(11):877-883. DOI:10.1016/j.jpplph.2014.03.003.
  77. Kato-Noguchi H, Moriyasu M, Ohno O, Suenaga K. Growth-limiting effects on various terrestrial plant species by an allelopathic substance, loliolide, from water hyacinth. *Aquatic Botany*. 2014;117:56-61. DOI:10.1016/j.aquabot.2014.05.001.
  78. Salam MA, Kato-Noguchi H. Evaluation of allelopathic potential of neem (*Azadirachta indica* A. Juss) against seed germination and seedling growth of different test plant species. *International Journal of Sustainable Agriculture*. 2010;2(2):20-25.
  79. Kato-Noguchi H, Nakamura K, Ohno O, Suenaga K, Okuda N. Asparagus decline: Autotoxicity and autotoxic compounds in asparagus rhizomes. *Journal of Plant Physiology*. 2017;213:23-29. DOI:10.1016/j.jpplph.2017.02.011.
  80. Charoenying P, Teerarak M, Laosinwattana C. An allelopathic substance isolated from *Zanthoxylum limonella* Alston fruit. *Scientia Horticulturae*. 2010;125(3):411-416. DOI:10.1016/j.scienta.2010.04.045.
  81. Sasikumar K, Vijayalakshmi C, Parthiban K. Allelopathic effects of *Eucalyptus* on black gram (*Phaseolus mungo* L.).
  82. Dadkhah A. Phytotoxic potential of sugar beet (*Beta vulgaris*) and *Eucalyptus camaldulensis* to control purslane (*Portulaca oleracea*) weed. *Allelopathy Journal*. 2013;63(1):46-51. DOI:10.1080/09064710.2012.711354.

83. Puig CG, Reigosa MJ, Valentão P, Andrade PB, Pedrol N. Unravelling the bioherbicide potential of *Eucalyptus globulus* Labill: Biochemistry and effects of its aqueous extract. PLoS ONE. 2018;13(2). DOI:10.1371/journal.pone.0192872.
84. Hasan M, Ahmad-Hamdani M, Rosli A, Hamdan H. Bioherbicides: An eco-friendly tool for sustainable weed management. *Plants*. 2021;10(6):1212. DOI:10.3390/plants10061212.
85. Mohsenzadeh S. Allelopathic potential of medicinal plants: Focus on Aloe, Ajwain and Rohida Tree.
86. Paton A, Putievsky E. Taxonomic problems and cytotoxic relationships between and within varieties of *Ocimum basilicum* and related species (Labiatae). *Kew Bulletin*. 1996:509-524. DOI:10.2307/4117026.
87. Verma SK, Kumar S, Pandey V, Verma RK, Patra DD. Phytotoxic effects of sweet basil (*Ocimum basilicum* L.) extracts on germination and seedling growth of commercial crop plants. *European Journal of Experimental Biology*. 2012;2(6):2310-2316.
88. Bączek K, Kosakowska O, Gniewosz M, Gientka I, Węglarz Z. Sweet basil (*Ocimum basilicum* L.) productivity and raw material quality from organic cultivation. *Agronomy*. 2019;9(6):279. DOI:10.3390/agronomy9060279.
89. Prakash P, Gupta N. Therapeutic uses of *Ocimum sanctum* Linn (Tulsi) with a note on eugenol and its pharmacological actions: a short review. *Indian Journal of Physiology and Pharmacology*. 2005;49(2):125.
90. Walton AA. *Ocimum basilicum*: Taxonomy, cultivation, and uses. *Journal of Botanical Taxonomy, Cultivation & Uses*. 2021:1-139.
91. Purohit S, Pandya N. Allelopathic activity of *Ocimum sanctum* L. and *Tephrosia purpurea* (L.) Pers. leaf extracts on few common legumes and weeds. *Indian Journal of Research in Pharmacy Science*. 2013;3(1):5-9.
92. Hiltunen R, Holm Y. Essential oil of *Ocimum*. In: *Basil*. CRC Press; 1999. p. 83-116.
93. Kamel E, Eid SD, Elian H. Allelochemicals effect of aqueous sweet basil (*Ocimum basilicum* L.) on weed control in peanut and cowpea crops. *Egyptian Journal of Chemistry*. 2022;65(5):153-164. DOI:10.21608/ejchem.2021.93101.4455.
94. Singh N, Thapar R. Allelopathic effects of *Croton bonplandianum* on *Parthenium hysterophorus*.
95. Knox J, Jaggi D, Paul M. Evaluation of allelopathic potential of selected plant species on *Parthenium hysterophorus*. *Egyptian Journal of Botany*. 2010;12.
96. Dhima K, Vasilakoglou I, Garane V, Ritzoulis C, Lianopoulou V, Panou-Philotheou E. Competitiveness and essential oil phytotoxicity of seven annual aromatic plants. *Weed Science*. 2010;58(4):457-465. DOI:10.1614/WS-D-10-00031.1.
97. Tahir J. Weeds control in forest ecosystem by the allelopathic potential of black walnut (*Juglans nigra* L.). College of Agriculture, Duhok University, Duhok, Iraq.
98. Souza Filho A, Bayma J, Guilhon G, Zoghbi M. Potentially allelopathic activity of the essential oil of *Ocimum americanum*. *Planta Daninha*. 2009;27(3):499-505.
99. Khan RA, Khan F, Ahmed M, Shah AS, Khan NA, Khan MR. Phytotoxic and antibacterial assays of crude methanolic extract of *Mentha longifolia* (Linn.). *African Journal of Pharmacy and Pharmacology*. 2011;5(12):1530-1533. doi:10.5897/AJPP11.374.
100. Kumar P, Mishra S, Malik A, Satya S. Insecticidal properties of *Mentha* species: A review. *Industrial Crops and Products*. 2011;34(1):802-817. DOI:10.1016/j.indcrop.2011.02.019.
101. Puig CG, Álvarez-Iglesias L, Pardo-Muras M, Andrade PB, Pedrol N. *Mentha suaveolens* as allelopathic biomass for weed control: phenolics, organic acids, and volatile organic compounds profiles. *Agronomy*. 2023;13(11):2793. DOI:10.3390/agronomy13112793.
102. Mahdavia F, Saharkhiz MJ. Phytotoxic activity of essential oil and water extract of peppermint (*Mentha piperita* L. CV. Mitcham). *J Appl Res Med Aromat Plants*. 2015;2(4):146-153. DOI:10.1016/j.jarmap.2015.09.003.
103. Campiglia E, Mancinelli R, Cavalieri A, Caporali F. Use of essential oils of cinnamon, lavender, and peppermint for weed control. *Ital J Agron*. 2007;2(2):171-178. DOI:10.4081/ija.2007.171.

104. Petrova ST, Valcheva EG, Velcheva IG. A case study of allelopathic effect on weeds in wheat. *Environ Biol.* 2015;7(1).
105. Nagaraja T, Deshmukh S. Phytotoxic effect of *Andrographis paniculata* Nees on metabolism of *Parthenium hysterophorus* L. *J Bot.* 2009;2(2):165-167.
106. Azirak S, Karaman S. Allelopathic effect of some essential oils and components on the germination of weed species. *Allelopathy J.* 2008;58(1):88-92. DOI:10.1080/09064710701228353.
107. Isik D, Mennan H, Cam M, Tursun N, Arslan M. Allelopathic potential of some essential oil-bearing plant extracts on common lambsquarters (*Chenopodium album* L.). *Res Crop.* 2016;67(3):455-459.
108. Argyropoulos EI, Eleftherohorinos IG, Vokou D. In vitro evaluation of essential oils from Mediterranean aromatic plants of the Lamiaceae for weed control in tomato and cotton crops. *Weed Res.* 2008;48(2):100-109.
109. Mukerjee SK. A revision of the Labiateae of the Indian Empire. Agricultural Trade Development and Promotion Bureau; 1938.
110. Shah M, Prajapati M, Patel J, Modi K. *Leucas aspera*: A review. *Pharmacogn Rev.* 2010;4(8):85-87. DOI:10.4103/0973-7847.65330.
111. Srinivasan R, Ravali B, Suvarchala P, Honey A, Tejaswini A, Neeraja P. *Leucas aspera*—medicinal plant: A review.
112. Das SN, Patro VJ, Dinda SC. A review: Ethnobotanical survey of genus *Leucas*. *Pharmacogn Rev.* 2012;6(12):100. DOI:10.4103/0973-7847.99943.
113. Nirmala KA, Kanchana M. *Leucas aspera*—A review of its biological activity. *Sci Res Pharm.* 2018;9(1).
114. Islam A, Ohno O, Suenaga K, Kato-Noguchi H. Suaveolic acid: A potent phytotoxic substance of *Hyptis suaveolens*. *Sci World J.* 2014. DOI:10.1155/2014/425942.
115. Islam AM, Kato-Noguchi H. Isolation and characterization of allelopathic substance from *Leucas aspera*.
116. Thang Lam L, Kato-Noguchi H. Assessment of the allelopathic potential of *Leucas cephalotes* (Roth) Spreng. extracts on the seedling growth of six test plants. *Plant Omics J.* 2021. DOI:10.21475/POJ.14.01.21.p3396.
117. Ma L, Xie CM, Li J, Lou FC, Hu LH. Daturametelins H, I, and J: Three new withanolide glycosides from *Datura metel* L. *Chem Biodivers.* 2006;3(2):180-186. DOI:10.1002/cbdv.200690021.
118. Dabur R, Ali M, Singh H, Gupta J, Sharma G. A novel antifungal pyrrole derivative from *Datura metel* leaves. *Drug Dev Res.* 2004;59(7):568-570.
119. Sahai M, Manickam M, Gupta M, Srivastava A, Ray A. Characterisation of a cerebroside isolated from the leaves of *Datura metel*. *J Indian Chem Soc.* 1999;76(2):95-97.
120. Valverde P, Fornoni J, Núñez-Farfán J. Evolutionary ecology of *Datura stramonium*: Equal plant fitness benefits of growth and resistance against herbivory. *J Evol Biol.* 2003;16(1):127-137. DOI:10.1046/j.1420-9101.2003.00482.x.
121. Alexander J, Auðunsson GA, Benford D, Cockburn A, Cravedi JP, Dogliotti E, Fink-Gremmels J. Glucosinolates as undesirable substances in animal feed. *EFSA J.* 2008;6(1):590. DOI:10.2903/j.efsa.2008.590.
122. Oseni O, Olarinoye C, Amoo I. Studies on chemical compositions and functional properties of thorn apple (*Datura stramonium* L) Solanaceae. *Afr J Food Sci.* 2011;5(2):40-44.
123. Yang B, Wang Q, Xia Y, Feng W, Kuang H. Withanolide compounds from the flower of *Datura metel* L. *Helv Chim Acta.* 2007;90(8):1522-1528.
124. Lovett J, Potts WC. Primary effects of allelochemicals of *Datura stramonium* L. *Plant Soil.* 1987;98:137-144.
125. Sakadzo N, Pahlia I, Muzemu S, Mandumbu R, Makaza K. Herbicidal effects of *Datura stramonium* (L.) leaf extracts on *Amaranthus hybridus* (L.) and *Tagetes minuta* (L.). *Afr J Agric Res.* 2018;13(34):1754-1760. DOI:10.5897/AJAR2018.13225.
126. de Almeida JD, Freitas H. Exotic naturalized flora of continental Portugal—A reassessment. *Botanica Complutensis.* 2006;30:117.
127. Giardina G, Raimondo FM, Spadaro V. A catalogue of plants growing in Sicily: Herbarium Mediterraneum Panormitanum; 2007.
128. Andreu J, Vilà M, Hulme PE. An assessment of stakeholder perceptions and management of noxious alien plants in Spain. *Environ Manag.* 2009;43:1244-1255. DOI:10.1007/s00267-009-9280-1.

129. May F, Ash J. An assessment of the allelopathic potential of Eucalyptus. Aust J Bot. 1990;38(3):245-254. DOI:10.1071/BT9900245.
130. Al-Naib FA, Al-Mousawi AJKJ. Allelopathic effects of Eucalyptus microtheca: Identification and characterization of phenolic compounds. Sci Educ. 1976;3:83-87.
131. Muller CH, Muller WH, Haines BL. Volatile growth inhibitors produced by aromatic shrubs. Science. 1964;143(3605):471-473. DOI:10.1126/science.143.3605.471.
132. Putnam A. Allelopathic chemicals: can natural plant herbicides help control weeds?
133. Iqbal Z, Hussain I, Hussain A, Ashraf MY. Genetic variability to essential oil contents and composition in five species of Eucalyptus. Pak J Bot. 2004;35(5; SPI):843-852.
134. Babu RC, Kandasamy O. Allelopathic effect of Eucalyptus globulus Labill. on *Cyperus rotundus* L. and *Cynodon dactylon* L. Pers. J Agron Crop Sci. 1997;179(2):123-126. DOI:10.1111/j.1439-037X.1997.tb00507.x.
135. Daneshmandi MS, Azizi M. Allelopathic effect of Eucalyptus globulus Labill. on bermuda grass (*Cynodon dactylon* (L.) Pers.) germination and rhizome growth. Iran J Med Aromat Plants Res. 2009;25(3):333-346. DOI:10.22092/ijmapr.2009.7148.
136. Azizi M, Fujii Y. Allelopathic effect of some medicinal plant substances on seed germination of *Amaranthus retroflexus* and *Portulaca oleraceae*. Acta Hort. 2005;699:59-66. DOI:10.17660/ActaHortic.2006.699.5.
137. El-Rokiek K, Eid R. Allelopathic effects of Eucalyptus citriodora on amaryllis and associated grassy weed. Planta Daninha. 2009;27:887-899. DOI:10.1590/S0100-83582009000500002.
138. Inderjit, Weston LA, Duke SO. Challenges, achievements and opportunities in allelopathy research. J Plant Interact. 2005;1(2):69-81. DOI:10.1080/17429140600622535.
139. Zimdahl RL. Fundamentals of weed science. Academic Press; 2018.
140. Ferguson JJ, Rathinasabapathi B, Chase CA. Allelopathy: how plants suppress other plants: HS944/hs186. Environ Horticulture. 2013.
141. Vidal RA, Bauman TT. Fate of allelochemicals in the soil. Ciencia Rural. 1997;27:351-357. DOI:10.1590/S0103-84781997000200032.
142. Kobayashi K. Factors affecting phytotoxic activity of allelochemicals in soil. Weed Biol Manag. 2004;4(1):1-7. DOI:10.1111/j.1445-6664.2003.00112.x.
143. Khanh TD, Hong NH, Xuan TD, Chung IM. Paddy weed control by medicinal and leguminous plants from Southeast Asia. Crop Prot. 2005;24(5):421-431. DOI:10.1016/j.cropro.2004.09.020.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Peer-review history:**

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/124339>